

New U.S. Utility Patent Application

Title: NOVEL ANTI-ARRHYTHMIC AND HEART FAILURE DRUGS THAT  
TARGET THE LEAK IN THE RYANODINE RECEPTOR (RYR2)

Inventors: Andrew R. Marks, Donald W. Landry, Shi Xian Deng, and Zhen Zhuang  
Cheng

NOVEL ANTI-ARRHYTHMIC AND HEART FAILURE DRUGS THAT TARGET THE  
LEAK IN THE RYANODINE RECEPTOR (RYR2)

RELATED APPLICATIONS

[0001] This application is a continuation-in-part of U.S. Patent Application Serial No.  
5 10/680,988, filed on October 7, 2003, which is a continuation-in-part of U.S. Patent  
Application Serial No. 10/608,723, filed on June 26, 2003, which is a continuation-in-part of  
U.S. Patent Application Serial No. 10/288,606, filed on November 5, 2002, which is a  
continuation of U.S. Patent Application Serial No. 09/568,474, filed on May 10, 2000, now  
U.S. Patent 6,489,125 B1, issued on December 3, 2002, the contents of which are hereby  
10 incorporated by reference herein.

STATEMENT OF GOVERNMENT INTEREST

[0002] This invention was made with government support under NIH Grant No. PO1  
HL 67849-01. As such, the United States government has certain rights in this invention.

BACKGROUND OF THE INVENTION

15 [0003] Despite advances in treatment, congestive heart failure remains an important  
cause of mortality in Western countries. Heart failure affects 5 million individuals in the  
United States alone, and is characterized by a 5-year mortality rate of ~50% (Levy *et al.*,  
Long-term trends in the incidence of and survival with heart failure. *N. Engl. J. Med.*,  
347:1397-402, 2002). An important hallmark of heart failure is reduced myocardial  
20 contractility (Gwathmey *et al.*, Abnormal intracellular calcium handling in myocardium from  
patients with end-stage heart failure. *Circ. Res.*, 61:70-76, 1987).

[0004] In healthy heart muscle, and other striated muscle, calcium-release channels  
on the sarcoplasmic reticulum (SR), including ryanodine receptors (RyRs), facilitate coupling  
of the action potential to a muscle cell's contraction (*i.e.*, excitation-contraction (EC),  
25 coupling). Contraction is initiated when calcium ( $\text{Ca}^{2+}$ ) is released from the SR into the  
surrounding cytoplasm. In heart failure, contractile abnormalities result, in part, from  
alterations in the signaling cascade that allows the cardiac action potential to trigger  
contraction. In particular, in failing hearts, the amplitude of the whole-cell  $\text{Ca}^{2+}$  transient is  
decreased (Beuckelmann *et al.*, Intracellular calcium handling in isolated ventricular  
30 myocytes from patients with terminal heart failure. *Circ.*, 85:1046-55, 1992; Gomez *et al.*,  
Defective excitation-contraction coupling in experimental cardiac hypertrophy and heart  
failure. *Science*, 276:800-06, 1997), and the duration prolonged (Beuckelmann *et al.*,

Intracellular calcium handling in isolated ventricular myocytes from patients with terminal heart failure. *Circ.*, 85:1046-55, 1992).

[0005] Cardiac arrhythmia, a common feature of heart failure, results in many of the deaths associated with the disease. Atrial fibrillation (AF) is the most common cardiac arrhythmia in humans, and represents a major cause of morbidity and mortality (Chugh *et al.*, Epidemiology and natural history of atrial fibrillation: clinical implications. *J. Am. Coll. Cardiol.*, 37:371-78, 2001; Falk, R.H., Atrial fibrillation. *N. Engl. J. Med.*, 344:1067-78, 2001). Despite AF's clinical importance, the molecular mechanisms underlying this arrhythmia are poorly understood, and treatment options are limited.

10 [0006] It is well established that structural and electrical remodeling – including shortening of atrial refractoriness, loss of rate-related adaptation of refractoriness (Wijffels *et al.*, Atrial fibrillation begets atrial fibrillation: a study in awake chronically instrumented goats. *Circulation*, 92:1954-68, 1995; Morillo *et al.*, Chronic rapid atrial pacing: structural, functional, and electrophysiological characteristics of a new model of sustained atrial  
15 fibrillation. *Circulation*, 91:1588-95, 1995; Elvan *et al.*, Pacing-induced chronic atrial fibrillation impairs sinus node function in dogs: electrophysiological remodeling. *Circulation*, 94:2953-60, 1996; Gaspo *et al.*, Functional mechanisms underlying tachycardia-induced sustained atrial fibrillation in a chronic dog model. *Circulation*, 96:4027-35, 1997), and shortening of the wavelength of re-entrant wavelets – accompany sustained tachycardia  
20 (Rensma *et al.*, Length of excitation wave and susceptibility to reentrant atrial arrhythmias in normal conscious dogs. *Circ. Res.*, 62:395-410, 1988). This remodeling is likely important in the development, maintenance and progression of atrial fibrillation.

[0007] Previous studies suggest that calcium handling may play a role in electrical remodeling in atrial fibrillation (Sun *et al.*, Cellular mechanisms of atrial contractile  
25 dysfunction caused by sustained atrial tachycardia. *Circulation*, 98:719-27, 1998; Goette *et al.*, Electrical remodeling in atrial fibrillation: time course and mechanisms. *Circulation*, 94:2968-74, 1996; Daoud *et al.*, Effect of verapamil and procainamide on atrial fibrillation-induced electrical remodeling in humans. *Circulation*, 96:1542-50, 1997; Yu *et al.*, Tachycardia-induced change of atrial refractory period in humans: rate dependency and  
30 effects of antiarrhythmic drugs. *Circulation*, 97:2331-37, 1998; Leistad *et al.*, Atrial contractile dysfunction after short-term atrial fibrillation is reduced by verapamil but increased by BAY K8644. *Circulation*, 93:1747-54, 1996; Tieleman *et al.*, Verapamil

reduces tachycardia-induced electrical remodeling of the atria. *Circulation*, 95:1945-53, 1997). However, regulation of RyR2 during atrial fibrillations has not previously been reported.

[0008] Approximately 50% of all patients with heart disease die from fatal cardiac arrhythmias. In some cases, a ventricular arrhythmia in the heart may be rapidly fatal – a phenomenon referred to as "sudden cardiac death" (SCD). Fatal ventricular arrhythmias (and SCD) may also occur in young, otherwise-healthy individuals who are not known to have structural heart disease. In fact, ventricular arrhythmia is the most common cause of sudden death in otherwise-healthy individuals.

10 [0009] Catecholaminergic polymorphic ventricular tachycardia (CPVT) is an inherited disorder in individuals with structurally-normal hearts. It is characterized by stress-induced ventricular tachycardia – a lethal arrhythmia that may cause SCD. In subjects with CPVT, physical exertion and/or stress induce bidirectional and/or polymorphic ventricular tachycardias that lead to SCD in the absence of detectable structural heart disease (Laitinen *et al.*, Mutations of the cardiac ryanodine receptor (RyR2) gene in familial polymorphic ventricular tachycardia. *Circulation*, 103:485-90, 2001; Leenhardt *et al.*, Catecholaminergic polymorphic ventricular tachycardia in children: a 7-year follow-up of 21 patients. *Circulation*, 91:1512-19, 1995; Priori *et al.*, Clinical and molecular characterization of patients with catecholaminergic polymorphic ventricular tachycardia. *Circulation*, 106:69-74, 2002; Priori *et al.*, Mutations in the cardiac ryanodine receptor gene (hRyR2) underlie catecholaminergic polymorphic ventricular tachycardia. *Circulation*, 103:196-200, 2001; Swan *et al.*, Arrhythmic disorder mapped to chromosome 1q42-q43 causes malignant polymorphic ventricular tachycardia in structurally normal hearts. *J. Am. Coll. Cardiol.*, 34:2035-42, 1999).

25 [0010] CPVT is predominantly inherited in an autosomal-dominant fashion. Individuals with CPVT have ventricular arrhythmias when subjected to exercise, but do not develop arrhythmias at rest. Linkage studies and direct sequencing have identified mutations in the human RyR2 gene, on chromosome 1q42-q43, in individuals with CPVT (Laitinen *et al.*, Mutations of the cardiac ryanodine receptor (RyR2) gene in familial polymorphic ventricular tachycardia. *Circulation*, 103:485-90, 2001; Priori *et al.*, Mutations in the cardiac ryanodine receptor gene (hRyR2) underlie catecholaminergic polymorphic ventricular tachycardia. *Circulation*, 103:196-200, 2001; Swan *et al.*, Arrhythmic disorder mapped to chromosome

30

1q42-q43 causes malignant polymorphic ventricular tachycardia in structurally normal hearts. *J. Am. Coll. Cardiol.*, 34:2035-42, 1999).

[0011] There are three types of ryanodine receptors, all of which are highly-related  $\text{Ca}^{2+}$  channels: RyR1, RyR2, and RyR3. RyR1 is found in skeletal muscle, RyR2 is found in the heart, and RyR3 is located in the brain. The type 2 ryanodine receptor (RyR2) is the major  $\text{Ca}^{2+}$ -release channel required for EC coupling and muscle contraction in cardiac striated muscle.

[0012] RyR2 channels are packed into dense arrays in specialized regions of the SR that release intracellular stores of  $\text{Ca}^{2+}$ , and thereby trigger muscle contraction (Marx *et al.*, Coupled gating between individual skeletal muscle  $\text{Ca}^{2+}$  release channels (ryanodine receptors). *Science*, 281:818-21, 1998). During EC coupling, depolarization of the cardiac-muscle cell membrane, in phase zero of the action potential, activates voltage-gated  $\text{Ca}^{2+}$  channels. In turn,  $\text{Ca}^{2+}$  influx through these channels initiates  $\text{Ca}^{2+}$  release from the SR via RyR2, in a process known as  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release (Fabiato, A., Calcium-induced release of calcium from the cardiac sarcoplasmic reticulum. *Am. J. Physiol.*, 245:C1-C14, 1983; Nabauer *et al.*, Regulation of calcium release is gated by calcium current, not gating charge, in cardiac myocytes. *Science*, 244:800-03, 1989). The RyR2-mediated,  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release then activates the contractile proteins which are responsible for cardiac muscle contraction.

[0013] RyR2 is a protein complex comprising four 565,000-dalton RyR2 polypeptides in association with four 12,000-dalton FK506 binding proteins (FKBPs), specifically FKBP12.6 (calstabin). FKBPs are *cis-trans* peptidyl-prolyl isomerases that are widely expressed and serve a variety of cellular functions (Marks, A.R., Cellular functions of immunophilins. *Physiol. Rev.*, 76:631-49, 1996). FKBP12 proteins are tightly bound to, and regulate the function of, the skeletal ryanodine receptor, RyR1 (Brillantes *et al.*, Stabilization of calcium release channel (ryanodine receptor) function by FK506-binding protein. *Cell*, 77:513-23, 1994; Jayaraman *et al.*, FK506 binding protein associated with the calcium release channel (ryanodine receptor). *J. Biol. Chem.*, 267:9474-77, 1992); the cardiac ryanodine receptor, RyR2 (Kaftan *et al.*, Effects of rapamycin on ryanodine receptor/ $\text{Ca}^{2+}$ -release channels from cardiac muscle. *Circ. Res.*, 78:990-97, 1996); a related intracellular  $\text{Ca}^{2+}$ -release channel, known as the type 1 inositol 1,4,5-triphosphate receptor (IP3R1) (Cameron *et al.*, FKBP12 binds the inositol 1,4,5-trisphosphate receptor at leucine-proline

(1400-1401) and anchors calcineurin to this FK506-like domain. *J. Biol. Chem.*, 272:27582-88, 1997); and the type I transforming growth factor  $\beta$  (TGF $\beta$ ) receptor (T $\beta$ RI) (Chen *et al.*, Mechanism of TGF $\beta$  receptor inhibition by FKBP12. *EMBO J.*, 16:3866-76, 1997).

FKBP12.6 binds to the RyR2 channel (one molecule per RyR2 subunit), stabilizes RyR2-channel function (Brillantes *et al.*, Stabilization of calcium release channel (ryanodine receptor) function by FK506-binding protein. *Cell*, 77:513-23, 1994), and facilitates coupled gating between neighboring RyR2 channels (Marx *et al.*, Coupled gating between individual skeletal muscle Ca<sup>2+</sup> release channels (ryanodine receptors). *Science*, 281:818-21, 1998), thereby preventing aberrant activation of the channel during the resting phase of the cardiac cycle.

[0014] Phosphorylation of cardiac RyR2 by protein kinase (PKA) is an important part of the "fight or flight" response that increases cardiac EC coupling gain by augmenting the amount of Ca<sup>2+</sup> released for a given trigger (Marks, A.R., Cardiac intracellular calcium release channels: role in heart failure. *Circ. Res.*, 87:8-11, 2000). This signaling pathway provides a mechanism by which activation of the sympathetic nervous system, in response to stress, results in increased cardiac output required to meet the metabolic demands of the stress responses. Upon binding of catecholamines,  $\beta$ 1- and  $\beta$ 2-adrenergic receptors activate adenylyl cyclase *via* a stimulatory G-protein, G $\alpha_s$ . Adenylyl cyclase increases intracellular cAMP levels, which activate the cAMP-dependent PKA. PKA phosphorylation of RyR2 increases the open probability of the channel by dissociating calstabin2 (FKBP12.6) from the channel complex. This, in turn, increases the sensitivity of RyR2 to Ca<sup>2+</sup>-dependent activation (Hain *et al.*, Phosphorylation modulates the function of the calcium release channel of sarcoplasmic reticulum from cardiac muscle. *J. Biol. Chem.*, 270:2074-81, 1995; Valdivia *et al.*, Rapid adaptation of cardiac ryanodine receptors: modulation by Mg<sup>2+</sup> and phosphorylation. *Science*, 267:1997-2000, 1995; Marx *et al.*, PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell*, 101:365-76, 2000).

[0015] Failing hearts (*e.g.*, in patients with heart failure and in animal models of heart failure) are characterized by a maladaptive response that includes chronic hyperadrenergic stimulation (Bristow *et al.*, Decreased catecholamine sensitivity and beta-adrenergic-receptor density in failing human hearts. *N. Engl. J. Med.*, 307:205-11, 1982). The pathogenic significance of this stimulation in heart failure is supported by therapeutic strategies that

decrease beta-adrenergic stimulation and left ventricular myocardial wall stress, and potentially reverse ventricular remodeling (Barbone *et al.*, Comparison of right and left ventricular responses to left ventricular assist device support in patients with severe heart failure: a primary role of mechanical unloading underlying reverse remodeling. *Circulation*, 104:670-75, 2001; Eichhorn and Bristow, Medical therapy can improve the biological properties of the chronically failing heart. A new era in the treatment of heart failure. *Circulation*, 94:2285-96, 1996). In heart failure, chronic beta-adrenergic stimulation is associated with the activation of beta-adrenergic receptors in the heart, which, through coupling with G-proteins, activate adenylyl cyclase and thereby increase intracellular cAMP concentration. cAMP activates cAMP-dependent PKA, which has been shown to induce hyperphosphorylation of RyR2. Thus, chronic heart failure is a chronic hyperadrenergic state (Chidsey *et al.*, Augmentation of plasma norepinephrine response to exercise in patients with congestive heart failure. *N. Engl. J. Med.*, 267:650, 1962) which results in several pathologic consequences, including PKA hyperphosphorylation of RyR2 (Marx *et al.*, PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell*, 101:365-76, 2000).

[0016] The PKA hyperphosphorylation of RyR2 has been proposed as a factor contributing to depressed contractile function and arrhythmogenesis in heart failure (Marks *et al.*, Progression of heart failure: is protein kinase a hyperphosphorylation of the ryanodine receptor a contributing factor? *Circulation*, 105:272-75, 2002; Marx *et al.*, PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell*, 101:365-76, 2000). Consistent with this hypothesis, PKA hyperphosphorylation of RyR2 in failing hearts has been demonstrated *in vivo*, both in animal models and in patients with heart failure undergoing cardiac transplantation (Antos *et al.*, Dilated cardiomyopathy and sudden death resulting from constitutive activation of protein kinase A. *Circ. Res.*, 89:997-1004, 2001; Marx *et al.*, PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell*, 101:365-76, 2000; Ono *et al.*, Altered interaction of FKBP12.6 with ryanodine receptor as a cause of abnormal  $\text{Ca}^{2+}$  release in heart failure. *Cardiovasc. Res.*, 48:323-31, 2000; Reiken *et al.*, Beta-adrenergic receptor blockers restore cardiac calcium release channel (ryanodine receptor) structure and function in heart failure. *Circulation*, 104:2843-48, 2001; Semsarian *et al.*, The L-type calcium channel

inhibitor diltiazem prevents cardiomyopathy in a mouse model. *J. Clin. Invest.*, 109:1013-20, 2002; Yano *et al.*, Altered stoichiometry of FKBP12.6 versus ryanodine receptor as a cause of abnormal  $\text{Ca}^{2+}$  leak through ryanodine receptor in heart failure. *Circulation*, 102:2131-36, 2000).

- 5     **[0017]**         In failing hearts, the hyperphosphorylation of RyR2 by PKA induces the dissociation of the regulatory FKBP12.6 subunit from the RyR2 channel (Marx *et al.*, PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell*, 101:365-76, 2000). This causes marked changes in the biophysical properties of the RyR2 channel. Such changes are evidenced by
- 10    increased open probability ( $P_o$ ), due to an increased sensitivity to  $\text{Ca}^{2+}$ -dependent activation (Brillantes *et al.*, Stabilization of calcium release channel (ryanodine receptor) function by FK506-binding protein. *Cell*, 77:513-23, 1994; Kaftan *et al.*, Effects of rapamycin on ryanodine receptor /  $\text{Ca}^{2+}$ -release channels from cardiac muscle. *Circ. Res.*, 78:990-97, 1996); destabilization of the channel, resulting in subconductance states; and impaired coupled
- 15    gating of the channels, resulting in defective EC coupling and cardiac dysfunction (Marx *et al.*, Coupled gating between individual skeletal muscle  $\text{Ca}^{2+}$  release channels (ryanodine receptors). *Science*, 281:818-21, 1998). Thus, PKA-hyperphosphorylated RyR2 is very sensitive to low-level  $\text{Ca}^{2+}$  stimulation, and this manifests itself as an SR  $\text{Ca}^{2+}$  leak through the hyperphosphorylated channel.
- 20    **[0018]**         The maladaptive response to stress in heart failure results in depletion of FKBP12.6 from the channel macromolecular complex. This leads to a shift to the left in the sensitivity of RyR2 to  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release, resulting in channels that are more active at low-to-moderate  $[\text{Ca}^{2+}]$  (Marx *et al.*, PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell*,
- 25    101:365-76, 2000; Yamamoto *et al.*, Abnormal  $\text{Ca}^{2+}$  release from cardiac sarcoplasmic reticulum in tachycardia-induced heart failure. *Cardiovasc. Res.*, 44:146-55, 1999; Yano *et al.*, Altered stoichiometry of FKBP12.6 versus ryanodine receptor as a cause of abnormal  $\text{Ca}^{2+}$  leak through ryanodine receptor in heart failure. *Circulation*, 102:2131-36, 2000). Over time, the increased "leak" through RyR2 results in resetting of the SR  $\text{Ca}^{2+}$  content to a lower
- 30    level, which in turn reduces EC coupling gain and contributes to impaired systolic contractility (Marx *et al.*, PKA phosphorylation dissociates FKBP12.6 from the calcium



release channel (ryanodine receptor): defective regulation in failing hearts. *Cell*, 101:365-76, 2000).

[0019] Additionally, a subpopulation of RyR2 that are particularly "leaky" can release SR  $\text{Ca}^{2+}$  during the resting phase of the cardiac cycle, diastole. This results in depolarizations of the cardiomyocyte membrane known as delayed after-depolarizations (DADs), which are known to trigger fatal ventricular cardiac arrhythmias (Wehrens *et al.*, FKBP12.6 deficiency and defective calcium release channel (ryanodine receptor) function linked to exercise-induced sudden cardiac death. *Cell*, 113:829-40, 2003).

[0020] In structurally-normal hearts, a similar phenomenon may be at work.

Specifically, it is known that exercise and stress induce the release of catecholamines that activate beta-adrenergic receptors in the heart. Activation of the beta-adrenergic receptors leads to hyperphosphorylation of RyR2 channels. Evidence also suggests that the hyperphosphorylation of RyR2 resulting from beta-adrenergic-receptor activation renders mutated RyR2 channels more likely to open in the relaxation phase of the cardiac cycle, increasing the likelihood of arrhythmias.

[0021] Cardiac arrhythmias are known to be associated with SR  $\text{Ca}^{2+}$  leaks in structurally-normal hearts. In these cases, the most common mechanism for induction and maintenance of ventricular tachycardia is abnormal automaticity. One form of abnormal automaticity, known as triggered arrhythmia, is associated with aberrant release of SR  $\text{Ca}^{2+}$ , which initiates DADs (Fozzard, H.A., Afterdepolarizations and triggered activity. *Basic Res. Cardiol.*, 87:105-13, 1992; Wit and Rosen, Pathophysiologic mechanisms of cardiac arrhythmias. *Am. Heart J.*, 106:798-811, 1983). DADs are abnormal depolarizations in cardiomyocytes that occur after repolarization of a cardiac action potential. The molecular basis for the abnormal SR  $\text{Ca}^{2+}$  release that results in DADs has not been fully elucidated. However, DADs are known to be blocked by ryanodine, providing evidence that RyR2 may play a key role in the pathogenesis of this aberrant  $\text{Ca}^{2+}$  release (Marban *et al.*, Mechanisms of arrhythmogenic delayed and early afterdepolarizations in ferret ventricular muscle. *J. Clin. Invest.*, 78:1185-92, 1986; Song and Belardinelli, ATP promotes development of afterdepolarizations and triggered activity in cardiac myocytes. *Am. J. Physiol.*, 267:H2005-11, 1994).

[0022] In view of the foregoing, it is clear that leaks in RyR2 channels are associated with a number of pathological states – in both diseased hearts and structurally-normal hearts.

Accordingly, methods to repair the leaks in RyR2 could prevent heart failure, and fatal arrhythmias and fibrillations, in millions of patients.

[0023] JTV-519 (4-[3-(4-benzylpiperidin-1-yl)propionyl]-7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine monohydrochloride; also known as k201 or ICP-Calstan 100), a derivative of 1,4-benzothiazepine, is a new modulator of calcium-ion channels. In addition to regulating  $\text{Ca}^{2+}$  levels in myocardial cells, JTV-519 also modulates the  $\text{Na}^+$  current and the inward-rectifier  $\text{K}^+$  current in guinea pig ventricular cells, and inhibits the delayed-rectifier  $\text{K}^+$  current in guinea pig atrial cells. Studies have shown that JTV-519 has a strong cardioprotective effect against catecholamine-induced myocardial injury, myocardial-injury-induced myofibrillar overcontraction, and ischemia/reperfusion injury. In experimental myofibrillar overcontraction models, JTV-519 demonstrated greater cardioprotective effects than propranolol, verapamil, and diltiazem. Experimental data also suggest that JTV-519 effectively prevents ventricular ischemia/reperfusion by reducing the level of intracellular  $\text{Ca}^{2+}$  overload in animal models.

#### 15 SUMMARY OF THE INVENTION


[0024] The present invention is based upon the surprising discovery that RyR2 is a target for treating and preventing heart failure and cardiac arrhythmias, including atrial fibrillations and cardiac arrhythmias that cause exercise-induced sudden cardiac death (SCD). As described herein, the inventors made mutant RyR2 channels with 7 different CPVT mutations, and studied their functions. All 7 mutants had functional defects that resulted in channels that became leaky (a calcium leak) when stimulated during exercise. The inventors' study is the first to identify a mechanism by which the SR calcium leak causes DADs. Remarkably, the defect in the mutant CPVT channels made the channels look like the leaky channels in the hearts of patients with end-stage heart failure – a disorder characterized by a high incidence of fatal cardiac arrhythmias. Therefore, the inventors demonstrate herein that the mechanism for the VT in CPVT is the same as the mechanism for VT in heart failure.

[0025] The inventors also disclose herein that the drug JTV-519 (k201 or ICP-Calstan 100), a member of the 1,4 benzothiazepine family of compounds, repairs the leak in RyR2 channels. As the inventors show herein, JTV-519 enhances binding of FKBP12.6 to PKA-phosphorylated RyR2, and to mutant RyR2s that otherwise have reduced affinity for, or do not bind to, FKBP12.6. This action of JTV-519 fixes the leak in RyR2 that triggers fatal


-10-

cardiac arrhythmias (cardiac death) and that contributes to atrial/ventricular fibrillations and heart muscle dysfunction in heart failure.


**[0026]** Accordingly, in one aspect, the present invention provides novel 1,4-benzothiazepine intermediates and derivatives, as well as methods for synthesizing same, and methods for assaying for same. In certain embodiments, these novel 1,4-benzothiazepine intermediates and derivatives include:




wherein R = aryl, alkenyl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl;



wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl;




wherein R =  $CO(CH_2)_nXR'_2$ ,  $SO_2(CH_2)_nXR'_2$ , or  $SO_2NH(CH_2)_nXR'_2$ , and X = N or S, and n = 1, 2, or 3, and R' = alkyl or cycloalkyl; and wherein m = 1 or 2;



wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ ,  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl; and wherein X = NH or O; and

-11-



(e)

wherein  $R_1 = OR', SR', NR'$ , alkyl, or halide, at position 2, 3, 4, or 5 on the phenyl ring, and  $R' =$  alkyl, aryl, or H; wherein  $R_2 =$  H, alkyl, or aryl; and wherein  $R_3 =$  H, alkyl, or aryl.

Additional embodiments may include the following compounds: S7, S-20, S-25, S-27, and

5 S36 (set forth in Appendix A).

**[0027]** Also provided are uses of the novel 1,4-benzothiazepine compounds in methods for limiting or preventing a decrease in the level of RyR2-bound FKBP12.6 in a subject; in methods for treating or preventing heart failure, atrial fibrillation, or exercise-induced cardiac arrhythmia in a subject; and in methods for preventing exercise-induced sudden cardiac death in a subject.

10

**[0028]** In a further aspect, the present invention provides a method for identifying an agent that enhances binding of RyR2 and FKBP12.6, comprising the steps of: (a) obtaining or generating a source of RyR2; (b) exposing the RyR2 to FKBP12.6, in the presence of a candidate agent; and (c) determining if the agent enhances the binding of RyR2 and FKBP12.6. In certain embodiments, the RyR2 may be unphosphorylated, PKA-phosphorylated, or PKA-hyperphosphorylated. Also provided are an agent identified by this method, and uses of the agent in methods for limiting or preventing a decrease in the level of RyR2-bound FKBP12.6 in a subject; in methods for treating or preventing heart failure, atrial fibrillation, or exercise-induced cardiac arrhythmia in a subject; and in methods for preventing exercise-induced sudden cardiac death in a subject.

15

20

**[0029]** In still another aspect, the present invention provides a method for identifying an agent for enhancing the binding of RyR2 and FKBP12.6, comprising the steps of: (a) obtaining or generating a source of FKBP12.6; (b) exposing the FKBP12.6 to RyR2, in the presence of a candidate agent; and (c) determining if the agent enhances the binding of RyR2 and FKBP12.6. In certain embodiments, the RyR2 may be unphosphorylated, PKA-phosphorylated, or PKA-hyperphosphorylated. Also provided are an agent identified by this method, and uses of the agent in methods for limiting or preventing a decrease in the level of RyR2-bound FKBP12.6 in a subject; in methods for treating or preventing heart failure, atrial fibrillation, or exercise-induced cardiac arrhythmia in a subject; and in methods for preventing exercise-induced sudden cardiac death in a subject.

25

30

[0030] Additional aspects of the present invention will be apparent in view of the description which follows.

#### BRIEF DESCRIPTION OF THE FIGURES

[0031] FIG. 1 demonstrates that JTV-519 prevents exercise-induced ventricular  
5 arrhythmias in FKBP12.6<sup>+/-</sup> mice. (A) Representative ambulatory electrocardiograms of an untreated FKBP12.6<sup>+/-</sup> mouse, an FKBP12.6<sup>+/-</sup> mouse treated with JTV-519, and an FKBP12.6<sup>-/-</sup> mouse treated with JTV-519. There were no significant differences in heart rate, or in any of the measured ECG parameters. (B) upper tracing: Example of sustained polymorphic ventricular tachycardia, recorded in an untreated FKBP12.6<sup>+/-</sup> mouse subjected  
10 to exercise testing and injection with 1.0 mg/kg epinephrine. middle tracing: Electrocardiogram of a JTV-519-treated FKBP12.6<sup>+/-</sup> mouse following the same protocol; no arrhythmias were detected. bottom tracing: Exercise-induced ventricular tachycardia (VT) in an FKBP12.6<sup>-/-</sup> mouse treated with JTV-519. The dotted line represents 16.31 seconds of VT that are not shown in the figure. 'P' indicates a P-wave, which is indicative of sinus rhythm  
15 following ventricular tachycardia. (C) Bar graph showing quantification of sudden cardiac death (left), sustained ventricular tachycardias (>10 beats, middle), and non-sustained ventricular tachycardias (3-10 abnormal beats, right) in FKBP12.6<sup>+/-</sup> and FKBP12.6<sup>-/-</sup> mice, either treated or not treated with JTV-519, respectively. It should be noted that treatment with JTV-519 completely prevented exercise- and epinephrine-induced arrhythmias in  
20 FKBP12.6<sup>+/-</sup> mice treated with JTV-519 (n = 9), as compared with untreated FKBP12.6<sup>+/-</sup> mice (n = 10) or JTV-519-treated FKBP12.6<sup>-/-</sup> mice (n = 5), suggesting that JTV-519 prevents arrhythmias and sudden death in FKBP12.6<sup>+/-</sup> mice by rebinding FKBP12.6 to RyR2.

[0032] FIG. 2 shows that JTV-519 prevents exercise-induced sudden cardiac death  
25 (SCD) by increasing the affinity of FKBP12.6 for RyR2 in FKBP12.6<sup>+/-</sup> mice. (A-B) Cardiac ryanodine receptors (RyR2) were immunoprecipitated using RyR2-5029 antibody. Shown are immunoblots (A) and bar graphs (B) representing the quantified amounts of RyR2, PKA-phosphorylated RyR2 (RyR2-pSer<sup>2809</sup> antibody), and FKBP12.6 in wild-type (FKBP12.6<sup>+/+</sup>) mice, FKBP12.6<sup>+/-</sup> mice, and FKBP12.6<sup>-/-</sup> under resting conditions, and following exercise,  
30 either in the absence or presence of JTV-519, respectively. Under resting conditions, ~70% of FKBP12.6 is associated with RyR2 in FKBP12.6<sup>+/-</sup> mice. Following exercise testing, the amount of FKBP12.6 associated with the RyR2 complex was dramatically decreased in

-13-

FKBP12.6<sup>+/-</sup> mice, but this could be rescued by treatment with JTV-519. (C) RyR2 single channels were isolated from hearts obtained following exercise testing and epinephrine injection. Shown are channels from FKBP12.6<sup>+/-</sup> mice, with and without pre-treatment with JTV-519, and channels from FKBP12.6<sup>-/-</sup> mice following JTV-519 pre-treatment. It should be noted that RyR2-channel function was normalized in the exercised FKBP12.6<sup>+/-</sup> mouse treated with JTV-519. The representative single channel from an exercised FKBP12.6<sup>-/-</sup> mouse after JTV-519 treatment shows that FKBP12.6 in the heart is required for the action of JTV-519. The dotted lines represent incomplete channel openings, or 'subconductance' openings, and are indicative of FKBP12.6-depleted RyR2 channels. Tracings on the left represent 5.0 sec, while tracings on the right represent 500 msec. In the figure, Po = open probability; To = average open times; Tc = average closed times; and c = closed state of the channel. (D) Summary bar graph showing average open probabilities of single RyR2 channels (see above). JTV-519 dramatically reduces the open probability of RyR2 from FKBP12.6<sup>+/-</sup> mice following exercise testing at diastolic calcium concentrations (150 nM).

15 [0033] FIG. 3 illustrates JTV-519 normalizes RyR2-channel gating by increased FKBP12.6 binding affinity to PKA-phosphorylated RyR2 channels. (A, B) Canine cardiac SR membranes (A) and recombinantly-expressed RyR2 channels (B) were prepared as described previously (Kaftan *et al.*, Effects of rapamycin on ryanodine receptor/Ca<sup>(2+)</sup>-release channels from cardiac muscle. *Circ. Res.*, 78:990-97, 1996). (A) Ryanodine receptors (RyR2) were phosphorylated with PKA catalytic subunit (40 U; Sigma Chemical Co., St. Louis, MO), in the presence or absence of the PKA inhibitor, PKI<sub>5-24</sub>, in phosphorylation buffer (8 mM MgCl<sub>2</sub>, 10 mM EGTA, and 50 mM Tris/PIPES; pH 6.8). Samples were centrifuged at 100,000x g for 10 min, and washed three times in imidazole buffer (10 mM imidazole; pH 7). Recombinantly-expressed FKBP12.6 (final concentration = 250 nM) was added to the samples, in the absence or presence of different concentrations of JTV-519. After a 60-min incubation, samples were centrifuged at 100,000x g for 10 min, and washed twice in imidazole buffer. Samples were heated to 95°C, and size-fractionated using SDS-PAGE. Immunoblotting of the SR microsomes was performed, as previously described (Jayaraman *et al.*, FK506 binding protein associated with the calcium release channel (ryanodine receptor). *J. Biol. Chem.*, 267:9474-77, 1992), with anti-FKBP12.6 antibody (1:1,000) and anti-RyR2-5029 antibody (1:3,000). The figure demonstrates that JTV-519 enables FKBP12.6 to bind to: (A) PKA-phosphorylated RyR2 (partial binding at 100 nM;

-14-

complete binding at 1000 nM) or (B) RyR2-S2809D mutant channels, which are constitutively PKA-phosphorylated RyR2 channels. (C-E) Single-channel studies showing increased open probability of RyR2 following PKA phosphorylation (D), as compared with PKA phosphorylation in the presence of the specific PKA inhibitor, PKI<sub>5-24</sub> (C). Single-channel function was normalized in PKA-phosphorylated RyR2 incubated with FKBP12.6 in the presence of JTV-519 (E). Channel openings are upward, the dash indicates the level of full openings (4 pA), and the letter 'c' indicates the closed state. Channels are shown at compressed (5 sec, upper tracing) and expanded (500 msec, lower tracing) time scales, and recordings are at 0 mV. Amplitude histograms (right) revealed increased activity and subconductance openings in PKA-phosphorylated RyR2, but not following treatment with JTV-519 and FKBP12.6. (F) Normalized plot of open probability as a function of cytosolic [Ca<sup>2+</sup>]. Incubation of PKA-phosphorylated RyR2 with FKBP12.6 in the presence of JTV-519 shifted the Ca<sup>2+</sup>-dependence of RyR2 activation towards the right, making it similar to the Ca<sup>2+</sup>-dependence of unphosphorylated channels.

15 [0034] FIG. 4 demonstrates the experimental protocol used to test effects of the inventors' novel JTV-519-related compounds (disclosed herein) on hERG-channel current. Whole-cell patch-clamp experiments were carried out with physiological solutions at room temperature, in CHO cells transfected with hERG channel. Voltage-clamp protocols are shown in the lower panels. In vehicle, 0.1% DMSO in the external solution was applied with the same time-protocol as that shown in the upper panel.

20 [0035] FIG. 5 illustrates the effects of JTV-519 and the inventors' novel JTV-519-related compound, S36 (disclosed herein), on hERG-channel currents elicited by 80-mV depolarization. Representative hERG-channel currents (I(Kr)) were recorded from CHO cells before (open circle) and after (closed circle) application of 1  $\mu$ M JTV-519 (left panel) or 25 1  $\mu$ M JTV-S36 (right panel). The voltage-clamp protocol is shown below the current traces. Currents were elicited during 400-msec depolarization to +80 mV, from a holding potential of -90 mV. It should be noted that, upon the 400-msec depolarization (which mimics the human action potential duration (APD)), hERG channels pass very little outward current, because they rapidly inactivate. Tail currents marked by circles in current traces were 30 elicited by return of the membrane potential to -40 mV, in the recovery from inactivation through the open state. Because the tail current is a major contributor to control of the APD,

-15-

effects of the drugs were evaluated by tail currents at -40 mV: JTV-519 = 83% block; JTV-S36 = 39% block.

[0036] FIG. 6 shows effects of JTV-519, E4031, and the inventors' novel JTV-519-related compound, S36 (disclosed herein), on activation of hERG-channel currents (traces). Representative hERG-channel I-V relationships are shown before (control, left panels) and after (central panels) application of 0.1% DMSO (vehicle; upper central panel), 1  $\mu$ M JTV-519 (middle central panel), and 1  $\mu$ M JTV-S36 (lower central panel). The right panel shows that 5  $\mu$ M E4031 (a class III anti-arrhythmic drug known to block hERG channels) completely blocked hERG-channel currents. (Note the tail currents at -40 mV). The voltage-clamp protocol is set forth in FIG. 4, as an I-V relationship.

[0037] FIG. 7 demonstrates effects of JTV-519 and the inventors' novel JTV-519-related compound, S36 (disclosed herein), on activation of hERG-channel currents. The hERG-channel I-V relationships are shown for peak tail currents (activation) before (open squares) and after (closed squares) application of 0.1% DMSO (vehicle; upper panel, 1  $\mu$ M JTV-519 (lower left panel), and 1  $\mu$ M JTV-S36 (lower right panel). Washout of the drugs is depicted with open triangles. The voltage-clamp protocol is set forth in FIG. 4, as an I-V relationship. It should be noted that JTV-S36 did not block hERG currents at negative potentials (0 mV; 20 mV depolarization) showing voltage-dependent block of I(Kr).

#### DETAILED DESCRIPTION OF THE INVENTION

[0038] As discussed above, catecholaminergic polymorphic ventricular tachycardia (CPVT) is an inherited disorder in individuals with structurally-normal hearts. It is characterized by stress-induced ventricular tachycardia, a lethal arrhythmia that may cause sudden cardiac death (SCD). Mutations in RyR2 channels, located on the sarcoplasmic reticulum (SR), have been linked to CPVT. To determine the molecular mechanism underlying the fatal cardiac arrhythmias in CPVT, the inventors studied CPVT-associated mutant RyR2 channels (*e.g.*, S2246L, R2474S, N4104K, R4497C).

[0039] All individuals with CPVT have exercise-induced cardiac arrhythmias. The inventors previously showed that exercise-induced arrhythmias and sudden death (in patients with CPVT) result from a reduced affinity of FKBP12.6 for RyR2. Herein, the inventors have demonstrated that exercise activates RyR2 as a result of phosphorylation by adenosine 3', 5'-monophosphate (cAMP)-dependent protein kinase (PKA). Mutant RyR2 channels,



-16-

which had normal function in planar lipid bilayers under basal conditions, were more sensitive to activation by PKA phosphorylation – exhibiting increased activity (open probability) and prolonged open states, as compared with wild-type channels. In addition, PKA-phosphorylated mutant RyR2 channels were resistant to inhibition by  $Mg^{2+}$ , a physiological inhibitor of the channel, and showed reduced binding to FKBP12.6 (which stabilizes the channel in the closed state). These findings indicate that, during exercise, when the RyR2 are PKA-phosphorylated, the mutant CPVT channels are more likely to open in the relaxation phase of the cardiac cycle (diastole), increasing the likelihood of arrhythmias triggered by SR  $Ca^{2+}$  leak. Since heart failure is a leading cause of death worldwide, methods to repair the leak in RyR2 could prevent fatal arrhythmias in millions of patients.

[0040] The inventors have further demonstrated herein that JTV-519, a benzothiazepine derivative, prevents lethal ventricular arrhythmias in mice heterozygous for the FKBP12.6 gene. JTV-519 reduced the open probability of RyR2, isolated from FKBP12.6<sup>+/-</sup> mice that died following exercise, by increasing the affinity of FKBP12.6 for PKA-phosphorylated RyR2. Moreover, JTV-519 normalized gating of CPVT-associated mutant RyR2 channels by increasing FKBP12.6 binding affinity. These data indicate that JTV-519 may prevent fatal ventricular arrhythmias by increasing FKBP12.6-RyR2 binding affinity.

#### Novel Methods of Treatment and Prevention

[0041] In accordance with the foregoing, the present invention provides a method for limiting or preventing a decrease in the level of RyR2-bound FKBP12.6 in cells of a subject. As used herein, "FKBP12.6" includes both an "FKBP12.6 protein" and an "FKBP12.6 analogue". Unless otherwise indicated herein, "protein" shall include a protein, protein domain, polypeptide, or peptide, and any fragment thereof. An "FKBP12.6 analogue" is a functional variant of the FKBP12.6 protein, having FKBP12.6 biological activity, that has 60% or greater amino-acid-sequence homology with the FKBP12.6 protein. As further used herein, the term "FKBP12.6 biological activity" refers to the activity of a protein or peptide that demonstrates an ability to associate physically with, or bind with, unphosphorylated or non-hyperphosphorylated RyR2 (*i.e.*, binding of approximately two fold, or, more preferably, approximately five fold, above the background binding of a negative control), under the conditions of the assays described herein, although affinity may be different from that of FKBP12.6.

[0042] In addition, as used herein, "RyR2" includes both an "RyR2 protein" and an "RyR2 analogue". An "RyR2 analogue" is a functional variant of the RyR2 protein, having RyR2 biological activity, that has 60% or greater amino-acid-sequence homology with the RyR2 protein. As used herein, the term "RyR2 analogue" includes RyR1 – the skeletal-muscle isoform of RyR2. The RyR2 of the present invention may be unphosphorylated, phosphorylated (*e.g.*, by PKA), or hyperphosphorylated (*e.g.*, by PKA). As further used herein, the term "RyR2 biological activity" refers to the activity of a protein or peptide that demonstrates an ability to associate physically with, or bind with, FKBP12.6 (*i.e.*, binding of approximately two fold, or, more preferably, approximately five fold, above the background binding of a negative control), under the conditions of the assays described herein, although affinity may be different from that of RyR2.

[0043] As described above, the cardiac ryanodine receptor, RyR2, is a protein complex comprising four 565,000-dalton RyR2 proteins in association with four 12,000-dalton FKBP12.6 proteins. FK506 binding proteins (FKBPs) are *cis-trans* peptidyl-prolyl isomerases that are widely expressed, and serve a variety of cellular functions. FKBP12.6 protein is tightly bound to, and regulates the function of, RyR2. FKBP12.6 binds to the RyR2 channel, one molecule per RyR2 subunit, stabilizes RyR2-channel function, and facilitates coupled gating between neighboring RyR2 channels, thereby preventing aberrant activation of the channel during the resting phase of the cardiac cycle. Accordingly, as used herein, the term "RyR2-bound FKBP12.6" includes a molecule of an FKBP12.6 protein that is bound to an RyR2 protein subunit or a tetramer of FKBP12.6 that is in association with a tetramer of RyR2.

[0044] In accordance with the method of the present invention, a "decrease" in the level of RyR2-bound FKBP12.6 in cells of a subject refers to a detectable decrease, diminution, or reduction in the level of RyR2-bound FKBP12.6 in cells of the subject. Such a decrease is limited or prevented in cells of a subject when the decrease is in any way halted, hindered, impeded, obstructed, or reduced by the administration of JTV-519 (as described below), such that the level of RyR2-bound FKBP12.6 in cells of the subject is higher than it would otherwise be in the absence of JTV-519.

[0045] The level of RyR2-bound FKBP12.6 in a subject may be detected by standard assays and techniques, including those readily determined from the known art (*e.g.*, immunological techniques, hybridization analysis, immunoprecipitation, Western-blot

analysis, fluorescence imaging techniques, and/or radiation detection, etc.), as well as any assays and detection methods disclosed herein. For example, protein may be isolated and purified from cells of a subject using standard methods known in the art, including, without limitation, extraction from the cells (*e.g.*, with a detergent that solubilizes the protein) where  
5 necessary, followed by affinity purification on a column, chromatography (*e.g.*, FTLC and HPLC), immunoprecipitation (with an antibody), and precipitation (*e.g.*, with isopropanol and a reagent such as Trizol). Isolation and purification of the protein may be followed by electrophoresis (*e.g.*, on an SDS-polyacrylamide gel). A decrease in the level of RyR2-bound FKBP12.6 in a subject, or the limiting or prevention thereof, may be determined by  
10 comparing the amount of RyR2-bound FKBP12.6 detected prior to the administration of JTV-519 (in accordance with methods described below) with the amount detected a suitable time after administration of JTV-519.

**[0046]** In the method of the present invention, a decrease in the level of RyR2-bound FKBP12.6 in cells of a subject may be limited or prevented, for example, by inhibiting  
15 dissociation of FKBP12.6 and RyR2 in cells of the subject; by increasing binding between FKBP12.6 and RyR2 in cells of the subject; or by stabilizing the RyR2-FKBP12.6 complex in cells of a subject. As used herein, the term "inhibiting dissociation" includes blocking, decreasing, inhibiting, limiting, or preventing the physical dissociation or separation of an FKBP12.6 subunit from an RyR2 molecule in cells of the subject, and blocking, decreasing,  
20 inhibiting, limiting, or preventing the physical dissociation or separation of an RyR2 molecule from an FKBP12.6 subunit in cells of the subject. As further used herein, the term "increasing binding" includes enhancing, increasing, or improving the ability of phosphorylated RyR2 to associate physically with FKBP12.6 (*e.g.*, binding of approximately two fold, or, more preferably, approximately five fold, above the background binding of a  
25 negative control) in cells of the subject, and enhancing, increasing, or improving the ability of FKBP12.6 to associate physically with phosphorylated RyR2 (*e.g.*, binding of approximately two fold, or, more preferably, approximately five fold, above the background binding of a negative control) in cells of the subject. Additionally, in the method of the present invention, a decrease in the level of RyR2-bound FKBP12.6 in cells of a subject may be limited or  
30 prevented by directly decreasing the level of phosphorylated RyR2 in cells of the subject, or by indirectly decreasing the level of phosphorylated RyR2 in the cells (*e.g.*, by targeting an enzyme (such as PKA) or another endogenous molecule that regulates or modulates the

functions or levels of phosphorylated RyR2 in the cells). Preferably, the level of phosphorylated RyR2 in the cells is decreased by at least 10% in the method of the present invention. More preferably, the level of phosphorylated RyR2 is decreased by at least 20%.

[0047] In accordance with the method of the present invention, a decrease in the level of RyR2-bound FKBP12.6 is limited or prevented in a subject, particularly in cells of a subject. The subject of the present invention may be any animal, including amphibians, birds, fish, mammals, and marsupials, but is preferably a mammal (*e.g.*, a human; a domestic animal, such as a cat, dog, monkey, mouse, or rat; or a commercial animal, such as a cow or pig). Additionally, the subject of the present invention is a candidate for exercise-induced cardiac arrhythmia. Exercise-induced cardiac arrhythmia is a heart condition (*e.g.*, a ventricular fibrillation or ventricular tachycardia, including any that leads to sudden cardiac death) that develops during/after a subject has undergone physical exercise. A "candidate" for exercise-induced cardiac arrhythmia is a subject who is known to be, or is believed to be, or is suspected of being, at risk for developing cardiac arrhythmia during/after physical exercise. Examples of candidates for exercise-induced cardiac arrhythmia include, without limitation, an animal/person known to have catecholaminergic polymorphic ventricular tachycardia (CPVT); an animal/person suspected of having CPVT; and an animal/person who is known to be, or is believed to be, or is suspected of being, at risk for developing cardiac arrhythmia during/after physical exercise, and who is about to exercise, is currently exercising, or has just completed exercise. As discussed above, CPVT is an inherited disorder in individuals with structurally-normal hearts. It is characterized by stress-induced ventricular tachycardia – a lethal arrhythmia that may cause sudden cardiac death. In subjects with CPVT, physical exertion and/or stress induce bidirectional and/or polymorphic ventricular tachycardias that lead to sudden cardiac death (SCD) in the absence of detectable structural heart disease. Individuals with CPVT have ventricular arrhythmias when subjected to exercise, but do not develop arrhythmias at rest.

[0048] In the method of the present invention, the cells of a subject are preferably striated muscle cells. A striated muscle is a muscle in which the repeating units (sarcomeres) of the contractile myofibrils are arranged in registry throughout the cell, resulting in transverse or oblique striations that may be observed at the level of a light microscope. Examples of striated muscle cells include, without limitation, voluntary (skeletal) muscle cells and cardiac muscle cells. In a preferred embodiment, the cell used in the method of the

-20-

present invention is a human cardiac muscle cell. As used herein, the term "cardiac muscle cell" includes cardiac muscle fibers, such as those found in the myocardium of the heart. Cardiac muscle fibers are composed of chains of contiguous heart-muscle cells, or cardiomyocytes, joined end to end at intercalated disks. These disks possess two kinds of cell  
5 junctions: expanded desmosomes extending along their transverse portions, and gap junctions, the largest of which lie along their longitudinal portions.

[0049] In the method of the present invention, a decrease in the level of RyR2-bound FKBP12.6 is limited or prevented in cells of a subject by administering JTV-519 to the subject; this would also permit contact between cells of the subject and JTV-519. JTV-519  
10 (4-[3-(4-benzylpiperidin-1-yl)propionyl]-7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine monohydrochloride), also known as k201, is a derivative of 1,4-benzothiazepine, and a modulator of calcium-ion channels. In addition to regulating  $\text{Ca}^{2+}$  levels in myocardial cells, JTV-519 modulates the  $\text{Na}^{+}$  current and the inward-rectifier  $\text{K}^{+}$  current in guinea pig ventricular cells, and inhibits the delayed-rectifier  $\text{K}^{+}$  current in guinea pig atrial cells.  
15 FK506 and rapamycin are drugs that may be used to design other compounds that stabilize RyR2-FKBP12.6 binding in cells of a subject who is a candidate for exercise-induced cardiac arrhythmia. FK506 and rapamycin both dissociate FKBP12.6 from RyR2. It is possible to design and/or screen for compounds that are structurally related to these drugs, but have the opposite effects.

20 [0050] In the method of the present invention, JTV-519 may be administered to a subject by way of a therapeutic composition, comprising JTV-519 and a pharmaceutically-acceptable carrier. The pharmaceutically-acceptable carrier must be "acceptable" in the sense of being compatible with the other ingredients of the composition, and not deleterious to the recipient thereof. The pharmaceutically-acceptable carrier employed herein is selected from  
25 various organic or inorganic materials that are used as materials for pharmaceutical formulations, and which may be incorporated as analgesic agents, buffers, binders, disintegrants, diluents, emulsifiers, excipients, extenders, glidants, solubilizers, stabilizers, suspending agents, tonicity agents, vehicles, and viscosity-increasing agents. If necessary, pharmaceutical additives, such as antioxidants, aromatics, colorants, flavor-improving agents,  
30 preservatives, and sweeteners, may also be added. Examples of acceptable pharmaceutical carriers include carboxymethyl cellulose, crystalline cellulose, glycerin, gum arabic, lactose,

magnesium stearate, methyl cellulose, powders, saline, sodium alginate, sucrose, starch, talc, and water, among others.

[0051] The pharmaceutical formulations of the present invention may be prepared by methods well-known in the pharmaceutical arts. For example, the JTV-519 may be brought  
5 into association with a carrier or diluent, as a suspension or solution. Optionally, one or more accessory ingredients (*e.g.*, buffers, flavoring agents, surface active agents, and the like) also may be added. The choice of carrier will depend upon the route of administration.

[0052] JTV-519 may be administered to a subject by contacting target cells (*e.g.*, cardiac muscle cells) *in vivo* in the subject with the JTV-519. JTV-519 may be contacted  
10 with (*e.g.*, introduced into) cells of the subject using known techniques utilized for the introduction and administration of proteins, nucleic acids, and other drugs. Examples of methods for contacting the cells with (*i.e.*, treating the cells with) JTV-519 include, without limitation, absorption, electroporation, immersion, injection, introduction, liposome delivery, transfection, transfusion, vectors, and other drug-delivery vehicles and methods. When the  
15 target cells are localized to a particular portion of a subject, it may be desirable to introduce the JTV-519 directly to the cells, by injection or by some other means (*e.g.*, by introducing the JTV-519 into the blood or another body fluid). The target cells may be contained in heart tissue of a subject, and may be detected in heart tissue of the subject by standard detection methods readily determined from the known art, examples of which include, without  
20 limitation, immunological techniques (*e.g.*, immunohistochemical staining), fluorescence imaging techniques, and microscopic techniques.

[0053] Additionally, the JTV-519 of the present invention may be administered to a human or animal subject by known procedures, including, without limitation, oral administration, parenteral administration, and transdermal administration. Preferably, the  
25 JTV-519 is administered parenterally, by epifascial, intracapsular, intracranial, intracutaneous, intrathecal, intramuscular, intraorbital, intraperitoneal, intraspinal, intrasternal, intravascular, intravenous, parenchymatous, subcutaneous, or sublingual injection, or by way of catheter. In one embodiment, the agent is administered to the subject by way of targeted delivery to cardiac muscle cells *via* a catheter inserted into the subject's  
30 heart.

[0054] For oral administration, a JTV-519 formulation may be presented as capsules, tablets, powders, granules, or as a suspension. The formulation may have conventional

additives, such as lactose, mannitol, corn starch, or potato starch. The formulation also may be presented with binders, such as crystalline cellulose, cellulose derivatives, acacia, corn starch, or gelatins. Additionally, the formulation may be presented with disintegrators, such as corn starch, potato starch, or sodium carboxymethylcellulose. The formulation also may  
5 be presented with dibasic calcium phosphate anhydrous or sodium starch glycolate. Finally, the formulation may be presented with lubricants, such as talc or magnesium stearate.

[0055] For parenteral administration (*i.e.*, administration by injection through a route other than the alimentary canal), JTV-519 may be combined with a sterile aqueous solution that is preferably isotonic with the blood of the subject. Such a formulation may be prepared  
10 by dissolving a solid active ingredient in water containing physiologically-compatible substances, such as sodium chloride, glycine, and the like, and having a buffered pH compatible with physiological conditions, so as to produce an aqueous solution, then rendering said solution sterile. The formulation may be presented in unit or multi-dose containers, such as sealed ampoules or vials. The formulation may be delivered by any mode  
15 of injection, including, without limitation, epifascial, intracapsular, intracranial, intracutaneous, intrathecal, intramuscular, intraorbital, intraperitoneal, intraspinal, intrasternal, intravascular, intravenous, parenchymatous, subcutaneous, or sublingual, or by way of catheter into the subject's heart.

[0056] For transdermal administration, JTV-519 may be combined with skin  
20 penetration enhancers, such as propylene glycol, polyethylene glycol, isopropanol, ethanol, oleic acid, *N*-methylpyrrolidone, and the like, which increase the permeability of the skin to the JTV-519, and permit the JTV-519 to penetrate through the skin and into the bloodstream. The JTV-519/enhancer composition also may be further combined with a polymeric substance, such as ethylcellulose, hydroxypropyl cellulose, ethylene/vinylacetate, polyvinyl  
25 pyrrolidone, and the like, to provide the composition in gel form, which may be dissolved in a solvent, such as methylene chloride, evaporated to the desired viscosity, and then applied to backing material to provide a patch.

[0057] In accordance with the method of the present invention, JTV-519 may be administered to the subject (and JTV-519 may be contacted with cells of the subject) in an  
30 amount effective to limit or prevent a decrease in the level of RyR2-bound FKBP12.6 in the subject, particularly in cells of the subject. This amount may be readily determined by the skilled artisan, based upon known procedures, including analysis of titration curves

established *in vivo*, and methods and assays disclosed herein. A suitable amount of JTV-519 effective to limit or prevent a decrease in the level of RyR2-bound FKBP12.6 in the subject may range from about 5 mg/kg/day to about 20 mg/kg/day, and/or may be an amount sufficient to achieve plasma levels ranging from about 300 ng/ml to about 1000 ng/ml.

5 Preferably, the amount of JTV-519 ranges from about 10 mg/kg/day to about 20 mg/kg/day.

[0058] In one embodiment of the present invention, the subject has not yet developed exercise-induced cardiac arrhythmia. In this case, the amount of JTV-519 effective to limit or prevent a decrease in the level of RyR2-bound FKBP12.6 in the subject may be an amount of JTV-519 effective to prevent exercise-induced cardiac arrhythmia in the subject. Cardiac  
10 arrhythmia is a disturbance of the electrical activity of the heart that manifests as an abnormality in heart rate or heart rhythm. As used herein, an amount of JTV-519 "effective to prevent exercise-induced cardiac arrhythmia" includes an amount of JTV-519 effective to prevent the development of the clinical impairment or symptoms of the exercise-induced cardiac arrhythmia (*e.g.*, palpitations, fainting, ventricular fibrillation, ventricular  
15 tachycardia, and sudden cardiac death). The amount of JTV-519 effective to prevent exercise-induced cardiac arrhythmia in a subject will vary depending upon the particular factors of each case, including the type of exercise-induced cardiac arrhythmia, the subject's weight, the severity of the subject's condition, and the mode of administration of the JTV-519. This amount may be readily determined by the skilled artisan, based upon known  
20 procedures, including clinical trials, and methods disclosed herein. In a preferred embodiment, the amount of JTV-519 effective to prevent the exercise-induced cardiac arrhythmia is an amount of JTV-519 effective to prevent exercise-induced sudden cardiac death in the subject. In another preferred embodiment, the JTV-519 prevents exercise-induced cardiac arrhythmia and exercise-induced sudden cardiac death in the subject.

25 [0059] Because of its ability to stabilize RyR2-bound FKBP12.6, and maintain and restore balance in the context of dynamic PKA phosphorylation and dephosphorylation of RyR2, JTV-519 may also be useful in treating a subject who has already started to experience clinical symptoms of exercise-induced cardiac arrhythmia. If the symptoms of arrhythmia are observed in the subject early enough, JTV-519 might be effective in limiting or  
30 preventing a further decrease in the level of RyR2-bound FKBP12.6 in the subject.

[0060] Accordingly, in still another embodiment of the present invention, the subject has been exercising, or is currently exercising, and has developed exercise-induced cardiac



arrhythmia. In this case, the amount of JTV-519 effective to limit or prevent a decrease in the level of RyR2-bound FKBP12.6 in the subject may be an amount of JTV-519 effective to treat exercise-induced cardiac arrhythmia in the subject. As used herein, an amount of JTV-519 "effective to treat exercise-induced cardiac arrhythmia" includes an amount of JTV-519 effective to alleviate or ameliorate the clinical impairment or symptoms of the exercise-induced cardiac arrhythmia (*e.g.*, palpitations, fainting, ventricular fibrillation, ventricular tachycardia, and sudden cardiac death). The amount of JTV-519 effective to treat exercise-induced cardiac arrhythmia in a subject will vary depending upon the particular factors of each case, including the type of exercise-induced cardiac arrhythmia, the subject's weight, the severity of the subject's condition, and the mode of administration of the JTV-519. This amount may be readily determined by the skilled artisan, based upon known procedures, including clinical trials, and methods disclosed herein. In a preferred embodiment, the JTV-519 treats exercise-induced cardiac arrhythmia in the subject.

[0061] The present invention further provides a method for treating exercise-induced cardiac arrhythmia in a subject. The method comprises administering JTV-519 to the subject in an amount effective to treat exercise-induced cardiac arrhythmia in the subject. A suitable amount of JTV-519 effective to treat exercise-induced cardiac arrhythmia in the subject may range from about 5 mg/kg/day to about 20 mg/kg/day, and/or may be an amount sufficient to achieve plasma levels ranging from about 300 ng/ml to about 1000 ng/ml. The present invention also provides a method for preventing exercise-induced cardiac arrhythmia in a subject. The method comprises administering JTV-519 to the subject in an amount effective to prevent exercise-induced cardiac arrhythmia in the subject. A suitable amount of JTV-519 effective to prevent exercise-induced cardiac arrhythmia in the subject may range from about 5 mg/kg/day to about 20 mg/kg/day, and/or may be an amount sufficient to achieve plasma levels ranging from about 300 ng/ml to about 1000 ng/ml. Additionally, the present invention provides a method for preventing exercise-induced sudden cardiac death in a subject. The method comprises administering JTV-519 to the subject in an amount effective to prevent exercise-induced sudden cardiac death in the subject. A suitable amount of JTV-519 effective to prevent exercise-induced sudden cardiac death in the subject may range from about 5 mg/kg/day to about 20 mg/kg/day, and/or may be an amount sufficient to achieve plasma levels ranging from about 300 ng/ml to about 1000 ng/ml.

[0062] In various embodiments of the above-described methods, the exercise-induced cardiac arrhythmia in the subject is associated with VT. In preferred embodiments, the VT is CPVT. In other embodiments of these methods, the subject is a candidate for exercise-induced cardiac arrhythmia, including candidates for exercise-induced sudden cardiac death.

5 [0063] In view of the foregoing methods, the present invention also provides use of JTV-519 in a method for limiting or preventing a decrease in the level of RyR2-bound FKBP12.6 in a subject who is a candidate for exercise-induced cardiac arrhythmia. The present invention also provides use of JTV-519 in a method for treating or preventing exercise-induced cardiac arrhythmia in a subject. Furthermore, the present invention  
10 provides use of JTV-519 in a method for preventing exercise-induced sudden cardiac death in a subject.

[0064] As discussed above and presented herein, the inventors' data show that protein kinase A (PKA) phosphorylation of the cardiac ryanodine receptor, RyR2, on serine 2809 activates the channel by releasing the FK506 binding protein, FKBP12.6. In failing hearts  
15 (including human hearts and animal models of heart failure), RyR2 is PKA-hyperphosphorylated, resulting in defective channels that have decreased amounts of bound FKBP12.6, and have increased sensitivity to calcium-induced activation. The net result of these changes is that the RyR2 channels are "leaky". These channel leaks can result in a depletion of intracellular stores of calcium to such an extent that there is no longer enough  
20 calcium in the sarcoplasmic reticulum (SR) to provide a strong stimulus for muscle contraction. This results in weak contraction of heart muscle. As a second consequence of the channel leaks, RyR2 channels release calcium during the resting phase of the heart cycle known as "diastole". This release of calcium during diastole can trigger the fatal arrhythmias of the hearts (*e.g.*, ventricular tachycardia and ventricular fibrillation) that cause sudden  
25 cardiac death (SCD).

[0065] The inventors have also shown that treatment of heart failure with a mechanical pumping device, referred to as a left ventricular assist device (LVAD), which puts the heart at rest and restores normalized function, is associated with a reduction in the PKA hyperphosphorylation of RyR2, and normalized function of the channel. Furthermore,  
30 the inventors have shown that treatment of dogs (who have pacing-induced heart failure) with beta-adrenergic blockers (beta blockers) reverses the PKA hyperphosphorylation of RyR2. Beta blockers inhibit the pathway that activates PKA. The conclusion which may be drawn

from the results of the inventors' work is that PKA phosphorylation of RyR2 increases the activity of the channel, resulting in the release of more calcium into the cell for a given trigger (activator) of the channel.

[0066] As further disclosed herein, the inventors have established that exercise-induced sudden cardiac death is associated with an increase in phosphorylation of RyR2 proteins (particularly CPVT-associated RyR2 mutant proteins) and a decrease in the level of RyR2-bound FKBP12.6. It is possible to use this mechanism to design effective drugs for preventing exercise-induced sudden cardiac death. A candidate agent having the ability to limit or prevent a decrease in the level of RyR2-bound FKBP12.6 may, as a consequence of this limiting or preventive activity, have an effect on an RyR2-associated biological event, thereby preventing exercise-induced sudden cardiac death.

[0067] Accordingly, the present invention further provides a method for identifying an agent for use in preventing exercise-induced sudden cardiac death. The method comprises the steps of: (a) obtaining or generating a culture of cells containing RyR2; (b) contacting the cells with a candidate agent; (c) exposing the cells to one or more conditions known to increase phosphorylation of RyR2 in cells; and (d) determining if the agent limits or prevents a decrease in the level of RyR2-bound FKBP12.6 in the cells. As used herein, an "agent" shall include a protein, polypeptide, peptide, nucleic acid (including DNA or RNA), antibody, Fab fragment, F(ab')<sub>2</sub> fragment, molecule, compound, antibiotic, drug, and any combination(s) thereof. An agent that limits or prevents a decrease in the level of RyR2-bound FKBP12.6 may be either natural or synthetic, and may be an agent reactive with (*i.e.*, an agent that has affinity for, binds to, or is directed against) RyR2 and/or FKBP12.6. As further used herein, a cell "containing RyR2" is a cell (preferably, a cardiac muscle cell) in which RyR2, or a derivative or homologue thereof, is naturally expressed or naturally occurs. Conditions known to increase phosphorylation of RyR2 in cells include, without limitation, PKA.

[0068] In the method of the present invention, cells may be contacted with a candidate agent by any of the standard methods of effecting contact between drugs/agents and cells, including any modes of introduction and administration described herein. The level of RyR2-bound FKBP12.6 in the cell may be measured or detected by known procedures, including any of the methods, molecular procedures, and assays known to one of

skill in the art or described herein. In one embodiment of the present invention, the agent limits or prevents a decrease in the level of RyR2-bound FKBP12.6 in the cells.

[0069] As disclosed herein, RyR2 has been implicated in a number of biological events in striated muscle cells. For example, it has been shown that RyR2 channels play an important role in EC coupling and contractility in cardiac muscle cells. Therefore, it is clear that preventive drugs designed to limit or prevent a decrease in the level of RyR2-bound FKBP12.6 in cells, particularly cardiac muscle cells, may be useful in the regulation of a number of RyR2-associated biological events, including EC coupling and contractility. Thus, once the candidate agent of the present invention has been screened, and has been determined to have a suitable limiting or preventive effect on decreasing levels of RyR2-bound FKBP12.6, it may be evaluated for its effect on EC coupling and contractility in cells, particularly cardiac muscle cells. It is expected that the preventive agent of the present invention will be useful for preventing exercise-induced sudden cardiac death.

[0070] Accordingly, the method of the present invention may further comprise the steps of: (e) contacting the candidate agent with a culture of cells containing RyR2; and (f) determining if the agent has an effect on an RyR2-associated biological event in the cells. As used herein, an "RyR2-associated biological event" includes a biochemical or physiological process in which RyR2 levels or activity have been implicated. As disclosed herein, examples of RyR2-associated biological events include, without limitation, EC coupling and contractility in cardiac muscle cells. According to this method of the present invention, a candidate agent may be contacted with one or more cells (preferably, cardiac muscle cells) *in vitro*. For example, a culture of the cells may be incubated with a preparation containing the candidate agent. The candidate agent's effect on an RyR2-associated biological event then may be assessed by any biological assays or methods known in the art, including immunoblotting, single-channel recordings and any others disclosed herein.

[0071] The present invention is further directed to an agent identified by the above-described identification method, as well as a pharmaceutical composition comprising the agent and a pharmaceutically-acceptable carrier. The agent may be useful for preventing exercise-induced sudden cardiac death in a subject, and for treating or preventing other RyR2-associated conditions. As used herein, an "RyR2-associated condition" is a condition, disease, or disorder in which RyR2 level or activity has been implicated, and includes an RyR2-associated biological event. The RyR2-associated condition may be treated or

-28-

prevented in the subject by administering to the subject an amount of the agent effective to treat or prevent the RyR2-associated condition in the subject. This amount may be readily determined by one skilled in the art. In one embodiment, the present invention provides a method for preventing exercise-induced sudden cardiac death in a subject, by administering  
5 the agent to the subject in an amount effective to prevent the exercise-induced sudden cardiac death in the subject.

[0072] The present invention also provides an *in vivo* method for identifying an agent for use in preventing exercise-induced sudden cardiac death. The method comprises the steps of: (a) obtaining or generating an animal containing RyR2; (b) administering a candidate  
10 agent to the animal; (c) exposing the animal to one or more conditions known to increase phosphorylation of RyR2 in cells; and (d) determining if the agent limits or prevents a decrease in the level of RyR2-bound FKBP12.6 in the animal. The method may further comprise the steps of: (e) administering the agent to an animal containing RyR2; and (f) determining if the agent has an effect on an RyR2-associated biological event in the animal.  
15 Also provided is an agent identified by this method; a pharmaceutical composition comprising this agent; and a method for preventing exercise-induced sudden cardiac death in a subject, by administering this agent to the subject in an amount effective to prevent the exercise-induced sudden cardiac death in the subject.

[0073] The inventors' work has demonstrated that compounds which block PKA  
20 activation would be expected to reduce the activation of the RyR2 channel, resulting in less release of calcium into the cell. Compounds that bind to the RyR2 channel at the FKBP12.6 binding site, but do not come off the channel when the channel is phosphorylated by PKA, would also be expected to decrease the activity of the channel in response to PKA activation or other triggers that activate the RyR2 channel. Such compounds would also result in less  
25 calcium release into the cell. In view of these findings, the present invention further provides additional assays for identifying agents that may be useful in preventing exercise-induced sudden cardiac death, in that they block or inhibit activation of RyR2.

[0074] By way of example, the diagnostic assays of the present invention may screen for the release of calcium into cells *via* the RyR2 channel, using calcium-sensitive fluorescent  
30 dyes (*e.g.*, Fluo-3, Fura-2, and the like). Cells may be loaded with the fluorescent dye of choice, then stimulated with RyR2 activators to determine whether or not compounds added to the cell reduce the calcium-dependent fluorescent signal (Brillantes *et al.*, Stabilization of

calcium release channel (ryanodine receptor) function by FK506-binding protein. *Cell*, 77:513-23, 1994; Gillo *et al.*, Calcium entry during induced differentiation in murine erythroleukemia cells. *Blood*, 81:783-92, 1993; Jayaraman *et al.*, Regulation of the inositol 1,4,5-trisphosphate receptor by tyrosine phosphorylation. *Science*, 272:1492-94, 1996).

- 5 Calcium-dependent fluorescent signals may be monitored with a photomultiplier tube, and analyzed with appropriate software, as previously described (Brillantes *et al.*, Stabilization of calcium release channel (ryanodine receptor) function by FK506-binding protein. *Cell*, 77:513-23, 1994; Gillo *et al.*, Calcium entry during induced differentiation in murine erythroleukemia cells. *Blood*, 81:783-92, 1993; Jayaraman *et al.*, Regulation of the inositol  
10 1,4,5-trisphosphate receptor by tyrosine phosphorylation. *Science*, 272:1492-94, 1996). This assay can easily be automated to screen large numbers of compounds using multiwell dishes.

- [0075] To identify compounds that inhibit the PKA-dependent activation of RyR2-mediated intracellular calcium release, an assay may involve the expression of recombinant RyR2 channels in a heterologous expression system, such as Sf9, HEK293, or CHO cells  
15 (Brillantes *et al.*, Stabilization of calcium release channel (ryanodine receptor) function by FK506-binding protein. *Cell*, 77:513-23, 1994). RyR2 could also be co-expressed with beta-adrenergic receptors. This would permit assessment of the effect of compounds on RyR2 activation, in response to addition of beta-adrenergic receptor agonists.

- [0076] The level of PKA phosphorylation of RyR2 which correlates with the degree  
20 of heart failure may also be assayed, and then used to determine the efficacy of compounds designed to block the PKA phosphorylation of the RyR2 channel. Such an assay may be based on the use of antibodies that are specific for the RyR2 protein. For example, the RyR2-channel protein may be immunoprecipitated, and then back-phosphorylated with PKA and [ $\gamma$ - $^{32}\text{P}$ ]-ATP. The amount of radioactive [ $^{32}\text{P}$ ] label that is transferred to the RyR2  
25 protein may be then measured using a phosphorimager (Marx *et al.*, PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell*, 101:365-76, 2000).

- [0077] Another assay of the present invention involves use of a phosphoepitope-specific antibody that detects RyR2 that is PKA phosphorylated on Ser 2809.  
30 Immunoblotting with such an antibody can be used to assess efficacy of therapy for heart failure and cardiac arrhythmias. Additionally, RyR2 S2809A and RyR2 S2809D knock-in mice may be used to assess efficacy of therapy for heart failure and cardiac arrhythmias.

Such mice further provide evidence that PKA hyperphosphorylation of RyR2 is a contributing factor in heart failure and cardiac arrhythmias, by showing that the RyR2 S2809A mutation inhibits heart failure and arrhythmias, and that the RyR2 S2809D mutation worsens heart failure and arrhythmias.

5 Novel Compounds, Methods of Synthesizing Same, and Uses of Same

[0078] 1,4-benzothiazepine derivatives, particularly 2,3,4,5-tetrahydro-1,4-benzothiazepine derivatives, are important building blocks in the preparation of biologically-active molecules, including JTV-519. The inventors have developed a novel process for preparing 1,4-benzothiazepine intermediate compounds, such as 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine. The inventors' process utilizes readily-available and inexpensive starting materials, and provides high yields of key 1,4-benzothiazepine intermediates.

[0079] In the early 1990s, Kaneko *et al.* (US Patent 5,416,066; WO 92/12148; JP4230681) disclosed that JTV-519 could be prepared by reacting 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (a 1,4-benzothiazepine intermediate) with acryloyl chloride, and then reacting the resulting product with 4-benzyl piperidine.

[0080] Two processes for the preparation of 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine and similar compounds have been previously reported in the literature. The first process, disclosed by Kaneko *et al.* (U.S. Patent No. 5,416,066), involved a synthetic route of six steps that started with 2,5-dihydroxybenzoic acid. In this process, 2,5-dihydroxybenzoic acid was selectively methylated with dimethyl sulfate. The resulting compound was then reacted with dimethylthiocarbamoyl chloride for 20 h, and then subjected to high temperature (270°C) for 9 h. The product of this step was refluxed with sodium methoxide in methanol for 20 h. The product of the reflux step was then reacted with 2-chloroethylamine, under basic conditions and at a high temperature, to produce a cyclized amide. The cyclized amide was reduced with LiAlH<sub>4</sub> to yield 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (a 1,4-benzothiazepine intermediate).


[0081] The second process for the preparation of 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine was disclosed by Hitoshi in a Japanese patent (JP 10045706). This process started with 2-bromo-5-methoxy benzaldehyde. The bromide was substituted with NaSMe, and the resulting product was oxidized with chlorine, followed by reflux in water, to yield disulfide dialdehyde. The dialdehyde was treated with 2-chloroethylamine, and the resulting

-31-

product was reduced with a reducing agent, such as  $\text{NaBH}_4$ . The resulting compound was cyclized to give 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine.


[0082] Initially, the inventors attempted to prepare the 1,4-benzothiazepine intermediate, 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine, using the methods described above. However, they found that the first process, described by Kaneko *et al.* (U.S. Patent No. 5,416,066), involved synthetic steps of high temperature and long reaction time. Additionally, the inventors discovered that the thio group in the third thiolated intermediate was easily oxidized by air to a disulfide compound, making it impossible to synthesize the subsequent cyclized product. The inventors also determined that the process described by Hitoshi (JP 10045706) involved  $\text{Cl}_2$ , and that another patented method for the preparation of the first intermediate, apart from the substitution of bromide with  $\text{NaSMe}$ , had to be used.

[0083] To overcome the foregoing problems, the inventors developed a novel process for making 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine from readily-available and inexpensive starting materials. The inventors' process simplifies isolation and purification steps, and can be used to prepare various 1,4-benzothiazepine intermediates, including 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine and other compounds having the general structure shown in formula:



This process may also be used to prepare JTV-519.

[0084] Accordingly, in view of the foregoing, the present invention provides a method for the synthesis of a compound having formula:




wherein R = OR', SR', NR', alkyl, or halide and R' = alkyl, aryl, or H, and wherein R can be at position 2, 3, 4, or 5, said method comprising the steps of:


(a) treating a compound having formula:



-32-




wherein R is as defined above, with a reducing agent, in the presence of an optional catalyst, to form a compound having formula:




5 wherein R is as defined above;

(b) treating the compound formed in step (a) with a diazotizing agent and a disulfide, to form a compound having formula:



wherein R is as defined above;


10 (c) treating the compound formed in step (b) with an activating agent and chloroethylamine, to form a compound having formula:



wherein R is as defined above;

(d) treating the compound formed in step (c) with a reducing agent and a base, to form a compound having formula:


15



wherein R is as defined above; and

(e) treating the compound formed in step (d) with a reducing agent, to form a compound having formula:


-33-



wherein R is as defined above.


[0085] In accordance with the method of the present invention, the reducing agent in step (a) may be  $H_2$ . Additionally, the diazotizing agent in step (b) may be  $NaNO_2$ , and the disulfide in step (b) may be  $Na_2S_2$ . Furthermore, the chloride in step (c) may be  $SOCl_2$ . The reducing agent in step (d) may be trimethylphosphine ( $PMe_3$ ), while the base in step (d) is triethyl amine. In another embodiment, the reducing agent in step (e) is  $LiAlH_4$ .

[0086] The present invention further provides a method for the synthesis of a compound of having formula:




wherein  $R = OR', SR', NR'$ , alkyl, or halide and  $R' =$  alkyl, aryl, or H, and wherein R can be at position 2, 3, 4, or 5, said method comprising the step of:


(a) treating a compound having formula:



wherein R is as defined above, with 3-bromopropionic chloride and a compound having formula:




to form a compound having formula:




wherein R is as defined above.

[0087] By way of example, a compound having the formula:

-34-



wherein R = OR', SR', NR', alkyl, or halide and R' = alkyl, aryl, or H, and wherein R can be at position 2, 3, 4, or 5, may be synthesized as follows:



R=OR', SR', NR', alkyl, halides; R'=alkyl, aryl, H  
R can be at positions 2, 3, 4, or 5

5

[0088] By way of example, and as shown in Example 7 and Scheme 1 below, 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine may be prepared from 2-nitro-5-methoxybenzoic acid as follows. The nitro group of 2-nitro-5-methoxybenzoic acid is reduced, using H<sub>2</sub> with Pd/C as a catalyst, to give 2-amino-5-methoxybenzoic acid. 2-amino-5-methoxybenzoic acid may be diazotized with NaNO<sub>2</sub>, and then treated with Na<sub>2</sub>S<sub>2</sub>, to provide a stable disulfide compound. Without further purification, the stable disulfide compound may be treated with SOCl<sub>2</sub>, and then reacted with 2-chloroethylamine, in the presence of Et<sub>3</sub>N, to give an amide. The amide compound may then be converted to a cyclized compound *via* a one-pot procedure, as follows. A reducing reagent (such as trimethylphosphine or triphenylphosphine) and a base (such as triethylamine) may be added to a solution of the amide compound in THF (tetrahydrofuran). The resulting reaction mixture may then be refluxed for 3 h. The reducing agent (trimethylphosphine or triphenylphosphine) cleaves the disulfide (S-S) to its monosulfide (-S), which, *in situ*, undergoes

15


-35-

intramolecular cyclization with the chloride to yield a cyclized amide. The cyclized amide may then be reduced with  $\text{LiAlH}_4$  to yield the 1,4-benzothiazepine intermediate, 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine. JTV-519 may then be prepared from 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine by reacting the 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine with 3-bromopropionic chloride, and then reacting the resulting compound with 4-benzyl piperidine.

[0089] The present invention further provides a composition, comprising radio-labeled JTV-519. Labeling of JTV-519 may be accomplished using one of a variety of different radioactive labels known in the art. The radioactive label of the present invention may be, for example, a radioisotope. The radioisotope may be any isotope that emits detectable radiation, including, without limitation,  $^{35}\text{S}$ ,  $^{125}\text{I}$ ,  $^3\text{H}$ , or  $^{14}\text{C}$ . Radioactivity emitted by the radioisotope can be detected by techniques well known in the art. For example, gamma emission from the radioisotope may be detected using gamma imaging techniques, particularly scintigraphic imaging.


[0090] By way of example, and as shown in Example 8 and Scheme 2 below, radio-labeled JTV-519 may be prepared as follows. JTV-519 may be demethylated at the phenyl ring using  $\text{BBr}_3$ . The resulting phenol compound may then be re-methylated with a radio-labeled methylating agent (such as  $^3\text{H}$ -dimethyl sulfate) in the presence of a base (such as  $\text{NaH}$ ) to provide  $^3\text{H}$ -labeled JTV-519.

[0091] The present invention further provides novel 1,4-benzothiazepine intermediates and derivatives, including 2,3,4,5-tetrahydro-1,4-benzothiazepenes that are similar to JTV-519. By way of example, the present invention provides compounds having the following formulas:




wherein  $\text{R} = \text{aryl}$ ,  $\text{alkenyl}$ ,  $\text{alkyl}$ ,  $-(\text{CH}_2)_n\text{NR}'_2$ , or  $-(\text{CH}_2)_n\text{SR}'$ , and  $n = 0, 1, 2$ , or  $3$ , and  $\text{R}' = \text{alkyl}$  or  $\text{cycloalkyl}$ ;

-36-




wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl;




5

wherein R =  $CO(CH_2)_nXR'_2$ ,  $SO_2(CH_2)_nXR'_2$ , or  $SO_2NH(CH_2)_nXR'_2$ , and X = N or S, and n = 1, 2, or 3, and R' = alkyl or cycloalkyl; and wherein m = 1 or 2; and



10 wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ ,  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl; and wherein X = NH or O. Also provided are additional 2, 3, 4, 5-tetrahydro-1,4-benzothiazepine compounds having formula:



15 wherein R<sub>1</sub> = OR', SR', NR', alkyl, or halide, at position 2, 3, 4, or 5 on the phenyl ring, and R' = alkyl, aryl, or H; wherein R<sub>2</sub> = H, alkyl, or aryl; and wherein R<sub>3</sub> = H, alkyl, or aryl.

[0092] Examples of the inventors' novel 1,4-benzothiazepine compounds include, without limitation, S7, S-20, S-25, S-27, and S36. Preferably, the compound is S36. Structures for S7, S-20, S-25, S-27, and S36 may be found in Appendix A, at the end of the present application. These and any other novel compounds of the present invention may be associated with a pharmaceutically-acceptable carrier, as described above, so as to form a pharmaceutical composition.

20 [0093] The inventors' novel 1,4-benzothiazepine compounds share functional characteristics with JTV-519. For example, like JTV-519 (mwt = 423), compound S36 (mwt = 267) regulates calcium channels. Indeed, S36 (a carboxylic acid) is approximately 10 times

-37-


more potent than JTV-519 in regulating calcium channels (data not shown). Unlike JTV-519, however, the inventors' novel compounds show weak blocking activity of hERGs.

[0094] The rapid delayed rectifier (I(Kr)) channel – a potassium channel – is important for repolarization of the cardiac action potential. hERG is the pore-forming subunit of the I(Kr) channel. Suppression of I(Kr) function – as a result of adverse drug effects and/or genetic defects in hERG – can lead to long-QT (LQT) syndrome, which carries increased risk of life-threatening arrhythmias. hERGs, then, are potassium-channel subunits that, when blocked, can cause cardiac arrhythmias and sudden cardiac death.

[0095] The inventors' compounds have significantly reduced blocking of hERG (I(Kr)) channels, when compared with JTV-519. As shown in FIGs. 4-7, for example, one of the inventors' compounds, S36, has hERG blocking activity that is approximately 5- to 10-fold lower than the hERG blocking activity of JTV-519. Because the inventors' compounds have weak hERG blocking activity, they are expected to be less toxic than JTV-519.


[0096] Based upon the foregoing, the inventors' novel compounds are more potent than JTV-519, and have reduced toxicity. Accordingly, it is believed that the inventors' novel compounds will be particularly useful in any of the above-described methods for limiting or preventing a decrease in the level of RyR2-bound FKBP12.6 in a subject, including a subject who is a candidate for heart failure, atrial fibrillation, or exercise-induced cardiac arrhythmia. It is also believed that the inventors' novel compounds will be particularly useful in methods for treating or preventing heart failure, atrial fibrillation, and exercise-induced cardiac arrhythmia in a subject, and in methods for preventing exercise-induced sudden cardiac death in a subject.

[0097] Accordingly, the present invention provides a method for limiting or preventing a decrease in the level of RyR2-bound FKBP12.6 in a subject, comprising administering to the subject an amount of agent effective to limit or prevent a decrease in the level of RyR2-bound FKBP12.6 in the subject, wherein the agent is selected from the group consisting of:




-38-

wherein R = aryl, alkenyl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl;



(b)


wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or  
5 cycloalkyl;



(c)

wherein R =  $CO(CH_2)_nXR'_2$ ,  $SO_2(CH_2)_nXR'_2$ , or  $SO_2NH(CH_2)_nXR'_2$ , and X = N or S, and n = 1, 2, or 3, and R' = alkyl or cycloalkyl; and wherein m = 1 or 2;


10



(d)

wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ ,  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or  
cycloalkyl; and wherein X = NH or O; and

15



(e)

wherein  $R_1 = OR'$ ,  $SR'$ ,  $NR'$ , alkyl, or halide, at position 2, 3, 4, or 5 on the phenyl ring, and  
R' = alkyl, aryl, or H; wherein  $R_2 = H$ , alkyl, or aryl; and wherein  $R_3 = H$ , alkyl, or aryl. As  
described above, the subject may be any animal, but is preferably a human. In one  
20 embodiment, the subject has catecholaminergic polymorphic ventricular tachycardia (CPVT).  
In another embodiment, the subject is a candidate for heart failure, atrial fibrillation, or  
exercise-induced cardiac arrhythmia. In still another embodiment, the agent is selected from

-39-


the group consisting of S4, S7, S-20, S-24, S-25, S-26, S-27, and S36. Structures for these agents may be found in Appendix A herein.

[0098] In accordance with the method of the present invention, the decrease in the level of RyR2-bound FKBP12.6 may be limited or prevented in the subject by decreasing the level of phosphorylated RyR2 in the subject. In one embodiment, the amount of the agent effective to limit or prevent a decrease in the level of RyR2-bound FKBP12.6 in the subject is an amount of the agent effective to treat or prevent heart failure, atrial fibrillation, and/or exercise-induced cardiac arrhythmia in the subject. In another embodiment, the amount of the agent effective to limit or prevent a decrease in the level of RyR2-bound FKBP12.6 in the subject is an amount of the agent effective to prevent exercise-induced sudden cardiac death in the subject.

[0099] In view of the foregoing, the present invention further provides a method for treating or preventing exercise-induced cardiac arrhythmia in a subject, comprising administering to the subject a novel 1,4-benzothiazepine compound, as disclosed herein, in an amount effective to treat or prevent exercise-induced cardiac arrhythmia in the subject. Similarly, the present invention provides a method for preventing exercise-induced sudden cardiac death in a subject, comprising administering to the subject a novel 1,4-benzothiazepine compound, as disclosed herein, in an amount effective to prevent exercise-induced sudden cardiac death in the subject. Additionally, the present invention provides a method for treating or preventing atrial fibrillation or heart failure in a subject, comprising administering to the subject a novel 1,4-benzothiazepine compound, as disclosed herein, in an amount effective to treat or prevent the atrial fibrillation or heart failure in the subject. In each of these methods, the novel 1,4-benzothiazepine compound may be selected from the group consisting of:

25


(a)



wherein R = aryl, alkenyl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl;




-40-




(b)

wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl;




(c)

wherein R =  $CO(CH_2)_nXR'_2$ ,  $SO_2(CH_2)_nXR'_2$ , or  $SO_2NH(CH_2)_nXR'_2$ , and X = N or S, and n = 1, 2, or 3, and R' = alkyl or cycloalkyl; and wherein m = 1 or 2;



(d)

wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ ,  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl; and wherein X = NH or O; and




(e)

wherein R<sub>1</sub> = OR', SR', NR', alkyl, or halide, at position 2, 3, 4, or 5 on the phenyl ring, and R' = alkyl, aryl, or H; wherein R<sub>2</sub> = H, alkyl, or aryl; and wherein R<sub>3</sub> = H, alkyl, or aryl.


**[00100]** The present invention further provides methods of synthesizing the novel 1,4-benzothiazepine compounds disclosed herein. For example, the present invention provides a method for the synthesis of a compound having formula:

-41-




wherein R = aryl, alkenyl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl, comprising the steps of:


- 5 (a) treating a compound having formula:



with a sulfonyl chloride compound (including any sulfonyl chloride derivative) and a base, to form a compound having the formula:




- 10 (b) optionally, treating the compound formed in step (a) with a primary or secondary amine, to form a compound having formula:



wherein R is as defined above. In one embodiment, the sulfonyl chloride compound in step (a) is selected from the group consisting of alkylsulfonyl chloride and arylsulfonyl chloride.


- 15 In another embodiment, the base in step (a) is  $Et_3N$ . In still another embodiment, the primary or secondary amine in step (b) is 4-benzylpiperidine.

**[00101]** The method of the invention may further comprise the step of oxidizing the compound having formula:



- 20 wherein R = aryl, alkenyl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl, with an oxidizing agent, to form a compound having formula:


-42-



wherein R is as defined above, and wherein  $m = 1$  or  $2$ . In one embodiment of the present invention, the oxidizing agent is hydrogen peroxide.


[00102] By way of example, and as shown in Example 9 and Scheme 3, the inventors

5 have developed a method of synthesizing compounds having the general structure:




wherein R = aryl, alkenyl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and  $n = 0, 1, 2$ , or  $3$ , and  $R' =$  alkyl or cycloalkyl. Novel compounds of this general structure may be prepared by reacting 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine with alkylsulfonyl chloride or arylsulfonyl chloride, in the presence of a base such as  $Et_3N$ . Additional reactions (*e.g.*, addition of 4-benzyl piperidine) may follow, to extend the side chain as desired. As Scheme 3 demonstrates, 2-chloroethanesulfonyl chloride (*e.g.*, 180 mg; 1.1 mM) and  $Et_3N$  (*e.g.*, 140 mg; 1.1 mM) may be added to 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (1) (*e.g.*, 194 mg; 1 mM) in  $CH_2Cl_2$  (*e.g.*, 20 ml), at  $0^\circ C$ . The mixture may then be stirred (*e.g.*, at  $0^\circ C$  for 2 h), and washed (*e.g.*, with  $H_2O$  and saturated  $NaHCO_3$  solution). Removal of the solvent will yield a crude product, which may be purified by chromatography on silica gel. The structure may be confirmed by NMR. Scheme 3 further shows that the resulting compound's side chain may be extended by reacting the compound (*e.g.*, 28 mg; 0.1 mM) with 4-benzyl piperidine (*e.g.*, 21 mg; 0.13 mM) in  $CH_2Cl_2$ . After the reaction goes to completion, the excess amine may be removed by a base scavenger (*e.g.*, 3-(2-succinic anhydride)propylfunctionalized silica gel, 0.5 g).

[00103] The present invention also provides a method for the synthesis of a compound of having formula:




-43-

wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl, comprising the step of treating a compound having formula:




- 5 with a sulfonyl chloride and a primary or secondary amine, in the presence of a base, to form a compound having the formula:




wherein R is as defined above. In one embodiment of the present invention, the base is  $Et_3N$ . In another embodiment, the primary or secondary amine is 1-piperonylpiperazine.

- [00104] The method of the present invention may further comprise the step of  
10 oxidizing the compound having formula:




wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl, to form a compound having formula:



- 15 wherein R is as defined above, and wherein m = 1 or 2. In one embodiment, the oxidizing agent is hydrogen peroxide.

[00105] By way of example, and as shown in Example 9 and Scheme 4, the inventors have developed a method of synthesizing compounds having the general structure:




- 20 wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl. Novel compounds of this general structure may be prepared by a one-pot reaction of 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine with sulfonyl chloride, in the presence of a base (*e.g.*,  $Et_3N$ ), followed by a primary or secondary amine. As Scheme 4 demonstrates, sulfonyl chloride (*e.g.*, 15.0 mg; 0.111 mM) and  $Et_3N$  (*e.g.*, 28.0 mg; 0.22 mM)

-44-


may be added to 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (*e.g.*, 19.4 mg; 0.1 mM) in  $\text{CH}_2\text{Cl}_2$  (*e.g.*, 20 ml), at  $0^\circ\text{C}$ . After stirring the mixture (*e.g.*, for 2 h at  $0^\circ\text{C}$ ), 1-piperonylpiperazine (*e.g.*, 27 mg; 0.12 mM) may be added. The mixture may be stirred for another 2 h, and then washed (*e.g.*, with  $\text{H}_2\text{O}$  and a saturated  $\text{NaHCO}_3$  solution). The excess amine may be removed by addition of a base scavenger (*e.g.*, 3-(2-succinic anhydride) propylfunctionalized silica gel, 0.2 g).

**[00106]** The present invention further provides a method for the synthesis of a compound of having formula:




10

wherein  $\text{R} = \text{CO}(\text{CH}_2)_n\text{XR}'_2$ ,  $\text{SO}_2(\text{CH}_2)_n\text{XR}'_2$ , or  $\text{SO}_2\text{NH}(\text{CH}_2)_n\text{XR}'_2$ , and  $\text{X} = \text{N}$  or  $\text{S}$ , and  $n = 1, 2$ , or  $3$ , and  $\text{R}' = \text{alkyl}$  or  $\text{cycloalkyl}$ ; and wherein  $m = 1$  or  $2$ , comprising the step of treating a compound having formula:




15 wherein R is as defined above, with an oxidizing agent, to form a compound having formula:



wherein R and m are as defined above. In one embodiment, the oxidizing agent is hydrogen peroxide. This method may also be used to oxidize JTV-519.

**[00107]** The present invention further provides a method for the synthesis of a compound of having formula:


20



wherein  $\text{R} = \text{CO}(\text{CH}_2)_n\text{XR}'_2$ ,  $\text{SO}_2(\text{CH}_2)_n\text{XR}'_2$ , or  $\text{SO}_2\text{NH}(\text{CH}_2)_n\text{XR}'_2$ , and  $\text{X} = \text{N}$  or  $\text{S}$ , and  $n = 1, 2$ , or  $3$ , and  $\text{R}' = \text{alkyl}$  or  $\text{cycloalkyl}$ ; and wherein  $m = 1$  or  $2$ , comprising the step of


25

treating a compound having formula:




-45-

with an oxidizing agent, to form a compound having formula:




wherein R and m are as defined above. In one embodiment, the oxidizing agent is hydrogen peroxide. This method may also be used to oxidize JTV-519.

- 5 **[00108]** By way of example, and as shown in Example 9 and Scheme 5, the inventors have developed a method of synthesizing compounds having the general structure:




- wherein R = CO(CH<sub>2</sub>)<sub>n</sub>XR'<sub>2</sub>, SO<sub>2</sub>(CH<sub>2</sub>)<sub>n</sub>XR'<sub>2</sub>, or SO<sub>2</sub>NH(CH<sub>2</sub>)<sub>n</sub>XR'<sub>2</sub>, and X = N or S, and n = 1, 2, or 3, and R' = alkyl or cycloalkyl; and wherein m = 1 or 2. Novel compounds of this  
 10 general structure may be prepared by oxidation of JTV-519, or one of the novel 1,4-benzothiazepine derivatives disclosed herein, with hydrogen peroxide. As Scheme 5 shows, the 1,4-benzothiazepine compound of interest (*e.g.*, 21 mg; 0.05 mM) in MeOH (*e.g.*, 5 ml) may be added to H<sub>2</sub>O<sub>2</sub> (*e.g.*, 0.1 ml, excess). The mixture may be stirred (*e.g.*, for 2 days), and the resulting product may be purified by chromatography on silica gel (*e.g.*,  
 15 CH<sub>2</sub>Cl<sub>2</sub>:MeOH = 10:1)

**[00109]** Additionally, the present invention provides a method for the synthesis of a compound having formula:




- wherein R=aryl, alkyl, -(CH<sub>2</sub>)<sub>n</sub>NR'<sub>2</sub>, or -(CH<sub>2</sub>)<sub>n</sub>SR', and n = 0, 1, 2, or 3, and R' = alkyl or  
 20 cycloalkyl; and wherein X = NH or O, comprising the step of treating a compound having formula:




with a carbonyl chloride compound, in the presence of a base, and with a primary or secondary amine or an alcohol, to form a compound having the formula:

-46-




wherein R and X are as defined above. In one embodiment, the carbonyl chloride compound is triphosgene. In another embodiment, the base is Et<sub>3</sub>N. In yet another embodiment, the primary or secondary amine is 4-benzylpiperidine.

- 5 **[00110]** By way of example, and as shown in Example 9 and Scheme 6, the inventors have developed a method of synthesizing compounds having the general structure:




- 10 wherein R = aryl, alkyl, -(CH<sub>2</sub>)<sub>n</sub>NR', -(CH<sub>2</sub>)<sub>n</sub>SR', and n = 0, 1, 2, or 3, and R' = alkyl or cycloalkyl; and wherein X = NH or O. Novel compounds of this general structure may be prepared by reacting 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine with triphosgene, in the presence of a base (*e.g.*, Et<sub>3</sub>N), followed by addition of a primary or secondary amine or an alcohol.

**[00111]** The present invention further provides a method for the synthesis of 2, 3, 4, 5-tetrahydro-1,4-benzothiazepine compounds having formula:




- 15 wherein R<sub>1</sub> = OR', SR', NR', alkyl, or halide, at position 2, 3, 4, or 5 on the phenyl ring, and R' = alkyl, aryl, or H; wherein R<sub>2</sub> = H, alkyl, or aryl; and wherein R<sub>3</sub> = H, alkyl, or aryl, comprising the steps of:

(a) treating a compound having formula:




- 20 wherein R<sub>1</sub> is as defined above, with a reducing agent, in the presence of an optional catalyst, to form a compound having formula:



wherein R<sub>1</sub> is as defined above;


-47-

(b) treating the compound formed in step (a) with a diazotizing agent and a disulfide, to form a compound having formula:




wherein R<sub>1</sub> is as defined above;

5 (c) treating the compound formed in step (b) with an activating agent and chloroethylamine, to form a compound having formula:




wherein R<sub>1</sub>, R<sub>2</sub>, and R<sub>3</sub> are as defined above;

(d) treating the compound formed in step (c) with a reducing agent and a base to  
10 form a compound having formula:



wherein R<sub>1</sub>, R<sub>2</sub>, and R<sub>3</sub> are as defined above; and

(e) treating the compound formed in step (d) with a reducing agent, to form a compound having formula:



15 wherein R<sub>1</sub>, R<sub>2</sub>, and R<sub>3</sub> are as defined above.

[00112] The present invention also provides novel assays for regular or high-throughput screening of biologically-active small molecules, based upon rebinding of FKBP12.6 and RyR2. In particular, the present invention provides a method for identifying an agent that  
20 enhances binding of RyR2 and FKBP12.6, comprising the steps of: (a) obtaining or



-48-

generating a source of RyR2; (b) exposing the RyR2 to FKBP12.6, in the presence of a candidate agent; and (c) determining if the agent enhances the binding of RyR2 and FKBP12.6. In one embodiment, the RyR2 is PKA-phosphorylated. In another embodiment, the RyR2 is PKA-hyperphosphorylated. In yet another embodiment, the RyR2 is  
5 unphosphorylated.

[00113] In the method of the present invention, the RyR2 is immobilized to a solid phase, such as a plate or beads. To facilitate detection of RyR2-FKBP12.6 binding, the FKBP12.6 may be radio-labeled (*e.g.*, with <sup>32</sup>S). Furthermore, enhanced binding of RyR2 and FKBP12.6 may be detected using an FKBP12.6-binding agent. In one embodiment, the  
10 FKBP12.6-binding agent is an anti-FKBP12.6 antibody. The present invention also provides an agent identified by this method, as well as uses of this agent in methods for limiting or preventing a decrease in the level of RyR2-bound FKBP12.6 in a subject; in methods for treating or preventing heart failure, atrial fibrillation, or exercise-induced cardiac arrhythmia in a subject; and in methods for preventing exercise-induced sudden cardiac death in a  
15 subject.

[00114] Additionally, the present invention provides a method for identifying an agent for enhancing the binding of RyR2 and FKBP12.6, comprising the steps of: (a) obtaining or generating a source of FKBP12.6; (b) exposing the FKBP12.6 to RyR2, in the presence of a candidate agent; and (c) determining if the agent enhances the binding of RyR2 and  
20 FKBP12.6. In one embodiment, the RyR2 is PKA-phosphorylated. In another embodiment, the RyR2 is PKA-hyperphosphorylated. In yet another embodiment, the RyR2 is unphosphorylated.

[00115] In the method of the present invention, the FKBP12.6 is immobilized to a solid phase, such as a plate or beads. To facilitate detection of RyR2-FKBP12.6 binding, the  
25 RyR2 may be radio-labeled (*e.g.*, with <sup>32</sup>P). Furthermore, enhanced binding of RyR2 and FKBP12.6 may be detected using an RyR2-binding agent. In one embodiment, the RyR2-binding agent is an anti-RyR2 antibody. The present invention also provides an agent identified by this method, as well as uses of this agent in methods for limiting or preventing a decrease in the level of RyR2-bound FKBP12.6 in a subject; in methods for treating or  
30 preventing heart failure, atrial fibrillation, or exercise-induced cardiac arrhythmia in a subject; and in methods for preventing exercise-induced sudden cardiac death in a subject.

-49-

[00116] By way of example, and as shown in Example 10 below, a highly-efficient assay for high-throughput screening for small molecules may be developed by immobilizing FKBP12.6 (*e.g.*, wild-type FKBP12.6 or a fusion protein, such as GST-FKBP12.6) onto a 96-well plate coated with glutathione, using standard procedures. PKA-phosphorylated ryanodine receptor type 2 (RyR2) may be loaded onto the FKBP12.6-coated plate, and incubated with JTV-519 analogues and other 1,4-benzothiazepene derivatives at various concentrations (10-100 nM) for 30 min. Thereafter, the plate may be washed to remove the unbound RyR2, and then incubated with anti-RyR2 antibody (*e.g.*, for 30 min). The plate may be washed again to remove unbound anti-RyR2 antibody, and then treated with fluorescent-labeled secondary antibody. The plate may be read by an automatic fluorescent plate reader for binding activity.

[00117] Alternatively, RyR2 may be PKA-phosphorylated in the presence of  $^{32}\text{P}$ -ATP. Radioactive PKA-phosphorylated RyR2 may be loaded onto an FKBP12.6-coated, 96-well plate, in the presence of JTV-519 analogues and other 1,4-benzothiazepene derivatives at various concentrations (10-100 nM) for 30 min. The plate may be washed to remove the unbound radiolabeled RyR2, and then read by an automatic plate reader. PKA-phosphorylated RyR2 also may be coated to the plate, and incubated with  $^{32}\text{S}$ -labeled FKBP12.6 in the presence of the analogues and derivatives.

[00118] The present invention is described in the following Examples, which are set forth to aid in the understanding of the invention, and should not be construed to limit in any way the scope of the invention as defined in the claims which follow thereafter.

## EXAMPLES

### EXAMPLE 1 – FKBP12.6-DEFICIENT MICE

[00119] FKBP12.6-deficient mice were generated, as previously described (Wehrens *et al.*, FKBP12.6 deficiency and defective calcium release channel (ryanodine receptor) function linked to exercise-induced sudden cardiac death. *Cell*, 113:829-40, 2003). Briefly, mouse genomic  $\lambda$ -phage clones for the murine orthologue of the human FK506 binding protein 12.6 (FKBP12.6) were isolated from a DBA/1lacJ library, using a full-length murine cDNA probe. The targeting vector was designed to delete exons 3 and 4, which contain the entire coding sequences for murine FKBP12.6 (Bennett *et al.*, Identification and characterization of the murine FK506 binding protein (FKBP) 12.6 gene. *Mamm. Genome*,

9:1069-71, 1998), by replacing 3.5 kb of murine genomic DNA with a PGK-neo selectable marker. A 5.0-kb 5' fragment and a 1.9-kb 3' fragment were cloned into pJNS2, a backbone vector with PGK-neo and PGK-TK cassettes. The DBA/lacJ embryonic stem (ES) cells were grown and transfected, using established protocols. Targeted ES cells were first screened by Southern analysis, and 5 positive ES cell lines were analyzed by PCR to confirm homologous recombination. Male chimeras were bred to DBA/1lacJ females, and germline offspring identified by brown coat color. Germline offspring were genotyped using 5' Southern analysis. Positive FKBP12.6<sup>+/-</sup> males and females were intercrossed, and offspring resulted in FKBP12.6<sup>-/-</sup> mice at approximately 25% frequency. FKBP12.6<sup>-/-</sup> mice were fertile.

- 10 [00120] All studies performed with FKBP12.6<sup>-/-</sup> mice used age- and sex-matched FKBP12.6<sup>+/+</sup> mice as controls. No differences were observed between FKBP12.6<sup>-/-</sup> mice raised on the following backgrounds: DBA/C57BL6 mixed, pure DBA, and pure C57BL6.

#### EXAMPLE 2 – TELEMETRY RECORDING AND EXERCISE TESTING IN MICE

- 15 [00121] FKBP12.6<sup>+/+</sup> and FKBP12.6<sup>-/-</sup> mice were maintained and studied according to protocols approved by the Institutional Animal Care and Use Committee of Columbia University. Mice were anaesthetized using 2.5% isoflurane inhalation anesthesia. ECG radiotelemetry recordings of ambulatory animals were obtained >7 days after intraperitoneal implantation (Data Sciences International, St. Paul, MN) (Wehrens *et al.*, FKBP12.6 deficiency and defective calcium release channel (ryanodine receptor) function linked to exercise-induced sudden cardiac death. *Cell*, 113:829-40, 2003). For stress tests, mice were exercised on an inclined treadmill until exhaustion, and then intraperitoneally injected with epinephrine (0.5-2.0 mg/kg) (Wehrens *et al.*, FKBP12.6 deficiency and defective calcium release channel (ryanodine receptor) function linked to exercise-induced sudden cardiac death. *Cell*, 113:829-40, 2003). Resting heart rates of ambulatory animals were averaged over 4 h.

#### EXAMPLE 3 – EXPRESSION OF WILD-TYPE AND RyR2-S2809D MUTANTS

- 30 [00122] Mutagenesis of the PKA target site on RyR2 (RyR2-S2809D) was performed, as previously described (Wehrens *et al.*, FKBP12.6 deficiency and defective calcium release channel (ryanodine receptor) function linked to exercise-induced sudden cardiac death. *Cell*, 113:829-40, 2003). HEK293 cells were co-transfected with 20 µg of RyR2 wild-type (WT) or mutant cDNA, and with 5 µg of FKBP12.6 cDNA, using Ca<sup>2+</sup> phosphate precipitation.

Vesicles containing RyR2 channels were prepared, as previously described (Wehrens *et al.*, FKBP12.6 deficiency and defective calcium release channel (ryanodine receptor) function linked to exercise-induced sudden cardiac death. *Cell*, 113:829-40, 2003).

#### EXAMPLE 4 – RyR2 PKA PHOSPHORYLATION AND FKBP12.6 BINDING

5 [00123] Cardiac SR membranes were prepared, as previously described (Marx *et al.*, PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell*, 101:365-76, 2000; Kaftan *et al.*, Effects of rapamycin on ryanodine receptor/Ca<sup>(2+)</sup>-release channels from cardiac muscle. *Circ. Res.*, 78:990-97, 1996). <sup>35</sup>S-labelled FKBP12.6 was generated using the TNT™ Quick Coupled  
10 Transcription/Translation system from Promega (Madison, WI). [<sup>3</sup>H] ryanodine binding was used to quantify RyR2 levels. 100 µg of microsomes were diluted in 100 µl of 10-mM imidazole buffer (pH 6.8), incubated with 250-nM (final concentration) [<sup>35</sup>S]-FKBP12.6 at 37°C for 60 min, then quenched with 500 µl of ice-cold imidazole buffer. Samples were centrifuged at 100,000 g for 10 min, and washed three times in imidazole buffer. The amount  
15 of bound [<sup>35</sup>S]-FKBP12.6 was determined by liquid scintillation counting of the pellet.

#### EXAMPLE 5 – IMMUNOBLOTS

[00124] Immunoblotting of microsomes (50 µg) was performed as described, with anti-FKBP12/12.6 (1:1,000), anti-RyR-5029 (1:3,000) (Jayaraman *et al.*, FK506 binding protein associated with the calcium release channel (ryanodine receptor). *J. Biol. Chem.*,  
20 267:9474-77, 1992), or anti-phosphoRyR2-P2809 (1:5,000) for 1 h at room temperature (Reiken *et al.*, Beta-blockers restore calcium release channel function and improve cardiac muscle performance in human heart failure. *Circulation*, 107:2459-66, 2003). The P2809-phosphoepitope-specific anti-RyR2 antibody is an affinity-purified polyclonal rabbit antibody, custom-made by Zymed Laboratories (San Francisco, CA) using the peptide,  
25 CRTRRI-(pS)-QTSQ, which corresponds to RyR2 PKA-phosphorylated at Ser<sup>2809</sup>. After incubation with HRP-labeled anti-rabbit IgG (1:5,000 dilution; Transduction Laboratories, Lexington, KY), the blots were developed using ECL (Amersham Pharmacia, Piscataway, NJ).

#### EXAMPLE 6 – SINGLE-CHANNEL RECORDINGS

30 [00125] Single-channel recordings of native RyR2 from mouse hearts, or recombinant RyR2, were acquired under voltage-clamp conditions at 0 mV, as previously described (Marx

*et al.*, PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell*, 101:365-76, 2000).

Symmetric solutions used for channel recordings were: *trans* compartment – HEPES, 250 mmol/L; Ba(OH)<sub>2</sub>, 53 mmol/L (in some experiments, Ba(OH)<sub>2</sub> was replaced by Ca(OH)<sub>2</sub>); pH 7.35; and *cis* compartment – HEPES, 250 mmol/L; Tris-base, 125 mmol/L; EGTA, 1.0 mmol/L; and CaCl<sub>2</sub>, 0.5 mmol/L; pH 7.35. Unless otherwise indicated, single-channels recordings were made in the presence of 150-nM [Ca<sup>2+</sup>] and 1.0-mM [Mg<sup>2+</sup>] in the *cis* compartment. Ryanodine (5 mM) was applied to the *cis* compartment to confirm identity of all channels. Data were analyzed from digitized current recordings using Fetchan software (Axon Instruments, Union City, CA). All data are expressed as mean ± SE. The unpaired Student's *t*-test was used for statistical comparison of mean values between experiments. A value of *p* < 0.05 was considered statistically significant.

[00126] The effects of JTV-519 on RyR2 channels are set forth in FIGs. 1-3 and Table 1 (below). As demonstrated in FIG. 3, the single-channel studies showed increased open probability of RyR2 following PKA phosphorylation (D), as compared to PKA phosphorylation in the presence of the specific PKA inhibitor, PKI<sub>5-24</sub> (C). Single-channel function was normalized in PKA-phosphorylated RyR2 incubated with FKBP12.6 in the presence of JTV-519 (E). Amplitude histograms (right) revealed increased activity and subconductance openings in PKA-phosphorylated RyR2, but not following treatment with JTV-519 and FKBP12.6. FIG. 3F shows that incubation of PKA-phosphorylated RyR2 with FKBP12.6, in the presence of JTV-519, shifted the Ca<sup>2+</sup>-dependence of RyR2 activation towards the right, making it similar to the Ca<sup>2+</sup>-dependence of unphosphorylated channels.

**Table 1. Ambulatory ECG data before, during exercise, and following exercise and injection with epinephrine.**

	SCL (ms)	HR (bpm)	PR (ms)	QRS (ms)	QT (ms)	QTc (ms)
Baseline						
FKBP12.6 <sup>+/+</sup>	104 ± 6	586 ± 36	32 ± 1.5	9.9 ± 0.4	30 ± 1.0	29 ± 0.6
FKBP12.6 <sup>+/+</sup> + JTV-519	99 ± 5	608 ± 32	33 ± 0.6	9.3 ± 0.3	32 ± 2.7	32 ± 1.9
FKBP12.6 <sup>-/-</sup> + JTV-519	116 ± 9	527 ± 43	33 ± 0.4	10.0 ± 0.3	33 ± 1.3	30 ± 1.1
Maximum exercise						
FKBP12.6 <sup>+/+</sup>	80 ± 2	752 ± 18	28 ± 0.7	8.7 ± 0.4	30 ± 1.7	33 ± 1.6
FKBP12.6 <sup>+/+</sup> + JTV-519	90 ± 7	676 ± 49	29 ± 1.8	9.6 ± 0.4	34 ± 2.0	36 ± 0.9
FKBP12.6 <sup>-/-</sup> + JTV-519	83 ± 3	729 ± 22	29 ± 2	9.3 ± 0.3	30 ± 1.2	33 ± 0.9

-53-

## Post-exercise epinephrine

FKBP12.6 <sup>+/-</sup>	94 ± 4	645 ± 28	35 ± 2.6	9.3 ± 0.4	33 ± 1.8	34 ± 1.9
FKBP12.6 <sup>+/-</sup> + JTV-519	102 ± 4	592 ± 21	37 ± 2.6	9.9 ± 0.6	32 ± 2.3	32 ± 1.7
FKBP12.6 <sup>-/-</sup> + JTV-519	103 ± 4	585 ± 20	35 ± 3.8	11.1 ± 0.5	36 ± 1.2	36 ± 1.3

Summary of ambulatory ECG data in FKBP12.6<sup>+/-</sup> mice treated with JTV-519 (n = 8) or control (n = 6), and FKBP12.6<sup>-/-</sup> mice treated with JTV-519 (n = 5). SCL = sinus cycle length; HR = heart rate; ms = millisecond; bpm = beats per minute; FKBP12.6<sup>+/-</sup> = FKBP12.6 heterozygous mice; FKBP12.6<sup>-/-</sup> = FKBP12.6 deficient mice

EXAMPLE 7 – SYNTHESIS OF 1,4-BENZOTHIAZEPINE  
INTERMEDIATE AND JTV-519


[00127] For the *in vivo* experiments, the inventors required a gram quantity of JTV-519. However, initial attempts to prepare this compound *via* the reported 1,4-benzothiazepine intermediate, 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (compound 6 in Scheme 1, below), were unsuccessful. The thio group of this intermediate is easily oxidized by air to a disulfide compound, which makes the synthesis of cyclized product (5) impossible. To overcome this problem, the inventors developed a novel process that starts with the readily-available and inexpensive 2-nitro-5-methoxybenzoic acid (1). This process is depicted in Scheme 1 below.

[00128] Reduction of the nitro group of compound (1), using H<sub>2</sub> with Pd/C as a catalyst, gave 2-amino-5-methoxybenzoic acid (2) in quantitative yield. Compound (2) was diazotized with NaNO<sub>2</sub>, and then treated with Na<sub>2</sub>S<sub>2</sub> to provide the stable disulfide compound (3) with 80% yield. Without further purification, the stable disulfide (3) was treated with SOCl<sub>2</sub>, and then reacted with 2-chloroethylamine, in the presence of Et<sub>3</sub>N, to give an amide (4) in 90% yield. Compound (4) was converted to cyclized compound (5) *via* a one-pot procedure by reflux with trimethylphosphine and Et<sub>3</sub>N in THF. The cyclized amide (5) was then reduced with LiAlH<sub>4</sub> to yield 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (6).

[00129] JTV-519 was prepared by reacting compound (6) with 3-bromopropionic chloride, and then reacting the resulting product with 4-benzyl piperidine. The structure of JTV-519 was established by <sup>1</sup>H NMR.


-54-

Scheme 1




#### EXAMPLE 8 – SYNTHESIS OF RADIO-LABELED JTV-519

[00130] The inventors' novel process for synthesizing radio-labeled JTV-519 is depicted in Scheme 2 below. To prepare radio-labeled JTV-519, JTV-519 was demethylated at the phenyl ring using BBr<sub>3</sub>, to give phenol compound (21). The phenol compound (21) was re-methylated with a radio-labeled methylating agent (<sup>3</sup>H-dimethyl sulfate) in the presence of a base (NaH) to provide <sup>3</sup>H-labeled JTV-519 (Scheme 2).



#### EXAMPLE 9 – NOVEL 1,4-BENZOTHAZEPINE DERIVATIVES AND METHODS FOR THEIR SYNTHESIS

[00131] The inventors also developed novel 1,4-benzothiazepine derivatives for use in treating and preventing cardiac arrhythmias. In particular, the inventors produced compounds having the following general structure:




-55-

wherein R = aryl, alkenyl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3; and wherein R' = alkyl or cycloalkyl. Novel compounds of this general structure were prepared by reacting 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine with alkylsulfonyl chloride or arylsulfonyl chloride, in the presence of a base such as  $Et_3N$ . Additional reactions (*e.g.*,  
 5 addition of 4-benzyl piperidine) may follow, to extend the side chain as desired. A representative synthesis of this general process is depicted in Scheme 3 below.


[00132] As Scheme 3 demonstrates, 2-chloroethanesulfonyl chloride (180 mg; 1.1 mM) and  $Et_3N$  (140 mg; 1.1 mM) were added to 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (1) (194 mg; 1 mM) in  $CH_2Cl_2$  (20 ml), at  $0^\circ C$ . The mixture was stirred at  
 10  $0^\circ C$  for 2 h, and washed with  $H_2O$  and saturated  $NaHCO_3$  solution. Removal of the solvent gave crude product (Ia), which was purified by chromatography on silica gel (petroleum ether : ethyl acetate = 3:1). The yield from this synthesis was 280 mg, or 95%. The structure was confirmed by NMR.

[00133] Scheme 3 further shows that the side chain of compound (Ia) was extended by  
 15 reacting compound (Ia) (28 mg; 0.1 mM) with 4-benzyl piperidine (21 mg; 0.13 mM) in  $CH_2Cl_2$ . After the reaction went to completion (by TLC), the excess amine was removed by a base scavenger (3-(2-succinic anhydride)propylfunctionalized silica gel, 0.5 g).  $^1H$ NMR and HPLC showed that the purity of product (Ib) was >98%.



25 Scheme 3

[00134] Additionally, the inventors produced compounds having the following general structure:






-56-


wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ , or  $-(CH_2)_nSR'$ , and n = 0, 1, 2, or 3; and wherein R' = alkyl or cycloalkyl. Novel compounds of this general structure were prepared by a one-pot reaction of 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (1) with sulfonyl chloride, in the presence of a base ( $Et_3N$ ), followed by a primary or secondary amine. A representative synthesis of this general process is depicted in Scheme 4 below.

[00135] As Scheme 4 demonstrates, sulfonyl chloride (15.0 mg; 0.111 mM) and  $Et_3N$  (28.0 mg; 0.22 mM) were added to 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (1) (19.4 mg; 0.1 mM) in  $CH_2Cl_2$  (20 ml), at  $0^\circ C$ . After stirring the mixture for 2 h at  $0^\circ C$ , 1-piperonylpiperazine (27 mg; 0.12 mM) was added. The mixture was stirred for another 2 h, and then washed with  $H_2O$  and a saturated  $NaHCO_3$  solution. The excess amine was removed by addition of a base scavenger (3-(2-succinic anhydride)propylfunctionalized silica gel, 0.2 g). The yield from this synthesis was 36 mg, or 77%.

Scheme 4




[00136] The inventors also produced compounds having the following general structure:



wherein R =  $CO(CH_2)_nXR'_2$ ,  $SO_2(CH_2)_nXR'_2$ , or  $SO_2NH(CH_2)_nXR'_2$ , and X = N or S, and n = 1, 2, or 3, and R' = alkyl or cycloalkyl; and wherein m = 1 or 2. Novel compounds of this general structure were prepared by oxidation of JTV-519, or one of the novel 1,4-benzothiazepine derivatives described above, with hydrogen peroxide. A representative synthesis of this general process is depicted in Scheme 5 below.


[00137] As Scheme 5 shows, compound (Ib) (21 mg; 0.05 mM) in MeOH (5 ml) was added to  $H_2O_2$  (0.1 ml, excess). The mixture was stirred for 2 days, and the product III was purified by chromatography on silica gel ( $CH_2Cl_2:MeOH = 10:1$ ). The yield from this synthesis was 19 mg, or 91%.

-57-



Scheme 5


[00138] Finally, the inventors produced compounds having the following general structure:



5

wherein R = aryl, alkyl,  $-(CH_2)_nNR'_2$ ,  $-(CH_2)_nSR'$ , and  $n = 0, 1, 2$ , or  $3$ , and  $R' =$  alkyl or cycloalkyl; and wherein  $X = NH$  or  $O$ . Novel compounds of this general structure were prepared by reacting 7-methoxy-2,3,4,5-tetrahydro-1,4-benzothiazepine (1) with triphosgene, in the presence of a base ( $Et_3N$ ), followed by addition of a primary or secondary amine or an alcohol. A representative synthesis of this general process is depicted in Scheme 6 below.

10



Scheme 6

#### EXAMPLE 10 – ASSAY FOR HIGH-THROUGHPUT SCREENING

[00139] The inventors have developed assays for screening biologically-active small molecules. These assays are based on rebinding of FKBP12 protein to RyR2.

15

[00140] A highly-efficient assay for high-throughput screening for small molecules may be developed by immobilization of FKBP12.6 (GST-fusion protein) onto a 96-well plate coated with glutathione. PKA-phosphorylated ryanodine receptor type 2 (RyR2) is loaded onto the FKBP12.6-coated plate, and incubated with JTV-519 analogues at various concentrations (10-100 nM) for 30 min. Thereafter, the plate is washed to remove the unbound RyR2, and then incubated with anti-RyR2 antibody for 30 min. The plate is again washed to remove unbound anti-RyR2 antibody, and then treated with fluorescent-labeled secondary antibody. The plate is read by an automatic fluorescent plate reader for binding activity.

20

-58-

[00141] In an alternative assay, RyR2 is PKA-phosphorylated in the presence of <sup>32</sup>P-ATP. Radioactive PKA-phosphorylated RyR2 is loaded onto an FKBP12.6-coated, 96-well plate, in the presence of JTV-519 analogues at various concentrations (10-100 nM) for 30 min. The plate is washed to remove the unbound radiolabeled RyR2, and then read by an automatic plate reader.

5

[00142] While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be appreciated by one skilled in the art, from a reading of the disclosure, that various changes in form and detail can be made without departing from the true scope of the invention in the appended claims.